

THE COTANGENT SPACE AT A MONOMIAL IDEAL OF THE HILBERT SCHEME OF POINTS OF AN AFFINE SPACE

MARK E. HUIBREGTSE

ABSTRACT. Let k be an algebraically closed field. We study the cotangent space of a point t corresponding to a monomial ideal $I \subseteq k[x_1, \dots, x_r]$ in the Hilbert scheme of n points of affine r -space (so $\dim_k(k[x_1, \dots, x_r]/I) = \text{colength of } I = n$). Since t lies in the closure of the locus corresponding to subschemes supported at n distinct points of $\mathbb{A}_k^r$, one knows that the k -dimension of the cotangent space is always $\geq rn$, and that t is nonsingular if and only if the dimension equals rn . We construct an explicit linearly independent set $\mathcal{S}$ of cotangent vectors of size rn , and then explore conditions on I under which $\mathcal{S}$ either is or is not a basis of the cotangent space. In particular, we give a condition on I sufficient for $\mathcal{S}$ to be a basis (equivalently, for t to be nonsingular) that holds for every monomial ideal in the case of $r = 2$ variables, and that characterizes such ideals when $r = 3$. We also give an easily-checked condition on I sufficient for $\mathcal{S}$ not to be a basis.

1. INTRODUCTION

1.1. Summary of results. Let k be an algebraically closed field of any characteristic, and

$$\mathbb{A}_k^r = \text{Spec}(k[x_1, \dots, x_r]) = \text{Spec}(k[\mathbf{x}])$$

the affine space of dimension r over k . The Hilbert scheme

$$\text{Hilb}_{\mathbb{A}_k^r}^n = \mathbf{H}^n$$

parameterizes the 0-dimensional closed subschemes

$$\text{Spec}(k[\mathbf{x}]/I) \subseteq \mathbb{A}_k^r$$

having length n , that is,

$$\dim_k(k[\mathbf{x}]/I) = \text{colength of } I = n.$$

In this paper we study the cotangent space of a point of $\mathbf{H}^n$ that corresponds to a monomial ideal I (that is, I is generated by monomials). If I is a monomial ideal of colength n , and we let

$$\beta = \{\text{monomials } m \mid m \notin I\},$$

Date: February 1, 2008.

1991 Mathematics Subject Classification. 14C05.

Key words and phrases. Hilbert scheme of points, affine space, monomial ideal, cotangent space.

then it is clear that β is a $\mathbf{k}$ -basis of the quotient $\mathbf{k}[\mathbf{x}]/I$; furthermore, β has the property that for monomials m_1, m_2 ,

$$m_1 \in \beta \text{ and } m_2 \mid m_1 \Rightarrow m_2 \in \beta;$$

we shall call any set of monomials β with this property a **basis set** of monomials. Let U_β denote the (affine) open subscheme of $\mathbf{H}^n$ whose $\mathbf{k}$ -points t are associated to ideals I_t such that β is a $\mathbf{k}$ -basis of the quotient $\mathbf{k}[\mathbf{x}]/I_t$. In particular, the monomial ideal $I = I_\beta$ that we started with has this property, so

$$t_\beta \in U_\beta, \text{ where } t_\beta \text{ is the point corresponding to } I_\beta = I_{t_\beta};$$

we can therefore identify the cotangent space of t_β with the $\mathbf{k}$ -vector space M/M^2 , where M is the maximal ideal of t_β in the coordinate ring R of U_β . Since t_β lies in the closure of the locus corresponding to closed subschemes supported at n distinct points of $\mathbb{A}_{\mathbf{k}}^r$, which is an (rn) -dimensional component of $\mathbf{H}^n$, we have that (Proposition 2.4.1)

- (1)
 - $\dim_{\mathbf{k}}(M/M^2) \geq rn$, and
 - t_β is nonsingular $\Leftrightarrow \dim_{\mathbf{k}}(M/M^2) = rn$.

Fortunately, we have a concrete description of R , based on the observation that for every point $t \in U_\beta$ and every monomial

$$x_1^{d_1} x_2^{d_2} \dots x_r^{d_r} = \mathbf{x}^{\mathbf{d}} \in I_\beta,$$

there is a unique polynomial

$$F_{\mathbf{d}}(t) = \mathbf{x}^{\mathbf{d}} - \sum_{\mathbf{x}^{\mathbf{j}} \in \beta} c_{\mathbf{j}}^{\mathbf{d}}(t) \cdot \mathbf{x}^{\mathbf{j}} \in I_t, \quad c_{\mathbf{j}}^{\mathbf{d}}(t) \in \mathbf{k},$$

since the quotient $\mathbf{k}[\mathbf{x}]/I_t$ has $\mathbf{k}$ -basis β . As t ranges over U_β , the coefficients $c_{\mathbf{j}}^{\mathbf{d}}(t)$ define functions $c_{\mathbf{j}}^{\mathbf{d}} \in R$ that generate R as a $\mathbf{k}$ -algebra; moreover, the relations among these functions can be completely described. Note in particular that the point t_β is the “origin” of U_β , the point at which all the functions $c_{\mathbf{j}}^{\mathbf{d}}$ vanish.

The maximal ideal $M \subseteq R$ that cuts out the point t_β is therefore generated by the functions $c_{\mathbf{j}}^{\mathbf{d}}$. Each of these functions can be viewed as an arrow pointing from $\mathbf{x}^{\mathbf{d}}$ to $\mathbf{x}^{\mathbf{j}}$ in the lattice of monomials obtained by identifying each monomial $\mathbf{x}^{\mathbf{b}}$ with its r -tuple $\mathbf{b}$ of exponents (for example, see Figures 1 and 2). It then turns out that translation of arrows (keeping the head inside β and the tail outside β) corresponds to equivalence modulo M^2 ; more precisely (see Section 3),

- If $c_{\mathbf{j}_1}^{\mathbf{d}_1}$ can be translated to $c_{\mathbf{j}_2}^{\mathbf{d}_2}$, then $c_{\mathbf{j}_1}^{\mathbf{d}_1} \equiv c_{\mathbf{j}_2}^{\mathbf{d}_2} \pmod{M^2}$.
- If $c_{\mathbf{j}_1}^{\mathbf{d}_1} \equiv 0 \pmod{M^2}$, then $c_{\mathbf{j}_1}^{\mathbf{d}_1}$ can be translated so that its head exits β across a hyperplane (x_i -degree = 0), and conversely.
- If $c_{\mathbf{j}_1}^{\mathbf{d}_1} \equiv c_{\mathbf{j}_2}^{\mathbf{d}_2} \not\equiv 0 \pmod{M^2}$, then $c_{\mathbf{j}_1}^{\mathbf{d}_1}$ can be translated to $c_{\mathbf{j}_2}^{\mathbf{d}_2}$.

Our first main result, Theorem 3.2.1, states that if S is a finite set of d arrows such that no member of S can be translated so that its head exits β across a hyperplane (x_i -degree = 0), and no two distinct members of S can be translated one to the other, then the $\mathbf{k}$ -span of S in M/M^2 has dimension d , and conversely; in this case, we say that S has **maximal rank** (mod M^2). Using this criterion, we construct in Theorem 4.1.3 a maximal rank (mod M^2) set $\mathcal{S}$ of rn arrows $c_j^{\mathbf{d}}$ of a special kind that we call **minimal standard** arrows: *minimal* means that the tail $\mathbf{x}^{\mathbf{d}}$ is a minimal generator of the monomial ideal I_β , and *standard* means that the vector $\mathbf{j} - \mathbf{d}$ has only one negative component. (Since $\mathbf{x}^{\mathbf{j}} \in \beta$ and $\mathbf{x}^{\mathbf{d}} \notin \beta$, the vector of any arrow must always have at least one negative component, to avoid the contradiction $\mathbf{x}^{\mathbf{d}} \mid \mathbf{x}^{\mathbf{j}}$.) Therefore, in view of (1),

$$t_\beta \text{ is nonsingular} \Leftrightarrow \mathcal{S} \text{ spans } M/M^2 \Leftrightarrow \mathcal{S} \text{ is a } \mathbf{k}\text{-basis of } M/M^2.$$

When any (and hence all) of these equivalent conditions holds, we say that β is a **smooth** basis set. (The set $\mathcal{S}$ is in general not unique.)

Remark 1.1.1. The idea of arrow translation and its connection to equivalence modulo M^2 is due to Haiman [6] (in the case of two variables); it is a pleasure to acknowledge the inspiration of that beautiful article.

The remainder of the paper explores conditions on β sufficient to imply that β either is or is not smooth. The basic result is Theorem 5.1.1, which in essence says that

$$(2) \quad \beta \text{ smooth} \Leftrightarrow \begin{cases} \bullet c_j^{\mathbf{d}} \text{ non-standard} \Rightarrow c_j^{\mathbf{d}} \equiv 0 \pmod{M^2}, \text{ and} \\ \bullet c_j^{\mathbf{d}'} \text{ standard and } \not\equiv 0 \pmod{M^2} \Rightarrow c_j^{\mathbf{d}'} \\ \quad \text{can be translated to an arrow in } \mathcal{S}. \end{cases}$$

In Section 5.2, we present a simple condition on β that ensures the existence of non-standard arrows that are *not* $\equiv 0$ modulo M^2 . One first checks that any minimal arrow $c_j^{\mathbf{d}}$ with head $\mathbf{x}^{\mathbf{j}}$ *maximal* in β (for the partial ordering defined by divisibility of monomials) cannot be translated except to itself (Lemma 5.2.1); we call such an arrow **rigid**. It is then clear that any non-standard rigid arrow will violate the first bulleted condition in (2), forcing β to be non-smooth (Corollary 5.2.2). Figure 1 illustrates such an arrow; a specific example is discussed in Section 5.3.

We go on to present three ways to construct smooth basis sets:

Thickening: In Section 6, we show that a basis set β in r variables that is the “thickening” (18) of a smooth basis set β_0 in $r - 1$ variables will be smooth (Theorem 6.4.1). As a consequence, we obtain that every **box** (a basis set that has the shape of a rectangular parallelepiped) is smooth (Proposition 6.5.3).

Box addition: In Section 9, we show that a basis set β that is obtained by adding a box to a smooth basis set in an appropriate way (see Subsection 9.1) is smooth (Theorem 9.3.1). Therefore, if we start with a box, and add a finite sequence of boxes to obtain a **compound box** (see Figure 6), we




FIGURE 1. A basis set β (the monomials within the boxes) in $r = 3$ variables with a non-standard rigid arrow c_j^b . The point $t_\beta \in \mathbf{H}^n$ is accordingly singular. The arrow is non-standard because the vector $\mathbf{j} - \mathbf{b}$ has 2 negative components: the first and the third.

obtain a smooth basis set (Corollary 9.4.1). In particular, *every* basis set in two variables is a compound box (see Figure 7) and therefore smooth, so we have obtained a variant proof of Haiman's result [6, Prop. 2.4, p. 209], using a lemma valid in arbitrary dimension. In addition, we prove in Section 10 that every smooth basis set in three variables is a compound box (Theorem 10.3.1).

Union of two boxes: In Section 11, we show that a basis set that is the union of two boxes is a smooth basis set (Theorem 11.3.2). We present this result to show that the characterization of smooth basis sets in three variables (as compound boxes) does not carry over to higher dimensions, because for $r \geq 4$ there are two-box unions that are not compound boxes (Example 11.4).

Sections 7 and 8 are devoted to technical preparations needed for Sections 9 and 10. Section 7 introduces the operation of *truncation* of a basis set (see Figure 4), and Section 8 establishes conditions under which a truncation of a smooth basis set will be smooth.

1.2. Table of contents.

1. Introduction.
 - 1.1 Summary of results.
 - 1.2 Table of contents.
2. The Hilbert scheme of points $\mathbf{H}^n$ and its open subschemes U_β .
 - 2.1 The Hilbert scheme $\mathbf{H}^n$.
 - 2.2 The open subschemes U_β .
 - 2.3 The coordinate ring of U_β .
 - 2.4 The locus of reduced subschemes meets U_β .
3. The cotangent space of the point $t_\beta \in \mathbf{H}^n$.
 - 3.1 Arrows, translation, and congruence modulo M^2 .

- 3.2 k -Linearly independent sets of arrows in M/M^2 .
- 4. A linearly independent set in M/M^2 of cardinality rn .
 - 4.1 Standard and minimal arrows.
 - 4.2 Advancement of minimal standard arrows.
 - 4.3 Shadow promotion.
 - 4.4 Iterated shadow promotion.
 - 4.5 Proof of Theorem 4.1.3
 - 4.6 Summary and terminology.
- 5. Consequences of Theorem 4.1.3.
 - 5.1 Necessary and sufficient conditions for $t_\beta \in \mathbf{H}^n$ to be nonsingular.
 - 5.2 A sufficient condition for $t_\beta \in \mathbf{H}^n$ to be singular.
 - 5.3 Example: $\beta = \{1, x_1, x_2, x_3\}$.
- 6. Thickening of basis sets.
 - 6.1 Definition of thickening.
 - 6.2 Minimal generators of I_β .
 - 6.3 A standard bunch $\mathcal{S}$ for β .
 - 6.4 Thickenings of smooth basis sets are smooth.
 - 6.5 Example: “Boxes” are smooth basis sets.
- 7. Truncation of basis sets.
 - 7.1 Definition of truncation.
 - 7.2 Minimal generators of I_{β_t} .
 - 7.3 Lifting arrows from β_t to β .
 - 7.4 Non-standard arrows on β and β_t .
- 8. Sufficient conditions for a truncation to be smooth.
 - 8.1 The additional hypothesis.
 - 8.2 x_j -sub-bunches of arrows for β and β_t .
 - 8.3 x_k -standard arrows of x_j -height $\geq h$.
 - 8.4 Linear independence of lifts of x_k -sub-bunches.
 - 8.5 x_k -sub-bunches of arrows for β and β_t , $x_k \neq x_j$.
 - 8.6 Main theorem on truncations.
- 9. Addition of boxes to basis sets.
 - 9.1 Definition of box addition.
 - 9.2 Minimal generators of $I_{\beta'}$.
 - 9.3 Main theorem on box additions.
 - 9.4 Compound boxes.
 - 9.5 Example: Basis sets in two variables.
 - 9.6 Example: The lexicographic point.
 - 9.7 Example: $\beta = \{1, x_1, x_2, x_1x_2, x_3\}$.
- 10. Smooth basis sets in three variables are compound boxes.

- 10.1 The main lemma.
- 10.2 Proof of Lemma 10.1.1.
- 10.3 The main theorem.
- 11. The union of two boxes.
 - 11.1 Notation.
 - 11.2 Minimal generators of I_β .
 - 11.3 Two-box unions are smooth basis sets.
 - 11.4 Example: $\beta = \{1, x_1, x_2, x_1x_2, x_3, x_4, x_3x_4\}$.

2. THE HILBERT SCHEME OF POINTS $\mathbf{H}^n$ AND ITS OPEN SUBSCHEMES U_β

In this section we briefly recall the definition and some properties of the Hilbert scheme of points $\mathbf{H}^n$, and the open subschemes U_β that form an open covering of $\mathbf{H}^n$. Much of our subsequent work is based on the explicit representation of the coordinate ring of U_β that is obtained in [9].

2.1. The Hilbert scheme $\mathbf{H}^n$. The scheme $\mathbf{H}^n$ can be defined functorially as the parameter scheme of a universal flat and proper family Z_n of zero-dimensional closed subschemes of $\mathbb{A}_k^r$ having length n . In other words, $Z_n \subseteq \mathbf{H}^n \times \mathbb{A}_k^r$ is a closed subscheme that is finite and flat of degree n over $\mathbf{H}^n$ and satisfies the following universal property:

(3) *Let T be a scheme over k . Then the set of maps $f: T \rightarrow \mathbf{H}^n$ is in natural bijective correspondence with the set of closed subschemes $Z_f \subseteq T \times \mathbb{A}_k^r$ that are finite and flat of degree n over T ; the bijection $f \mapsto Z_f$ is defined by $Z_f = T \times_{\mathbf{H}^n} Z_n$.*

In particular, the inclusion of the k -point $t \in \mathbf{H}^n$ corresponds to a unique closed subscheme

$$Z_t \subseteq t \times \mathbb{A}_k^r \approx \mathbb{A}_k^r;$$

the map $t \mapsto Z_t$ defines a bijection from the set of k -points of $\mathbf{H}^n$ to the set of 0-dimensional closed subschemes of length n , or, equivalently, to the set of ideals

$$I \subseteq k[x_1, \dots, x_r] = k[\mathbf{x}]$$

of colength n . We write I_t to denote the ideal corresponding to the k -point $t \in \mathbf{H}^n$.

The existence of $\mathbf{H}^n$ can be established as follows: It is the open subscheme of $\text{Hilb}_{\mathbb{P}_k^r}^n$ (a projective scheme the existence of which is a consequence of Grothendieck's general construction given in [4]) arising from the inclusion of $\mathbb{A}_k^r$ in $\mathbb{P}_k^r$ as a standard affine. Alternatively, $\mathbf{H}^n$ is a special case of the multigraded Hilbert scheme constructed by Haiman and Sturmfels [7]. Elementary constructions of $\mathbf{H}^n$ are given in [5] and [9].

2.2. The open subschemes U_β . Let β denote a nonempty set of n monomials in the indeterminates $x_1, \dots, x_r$ that satisfies the following property: if $m_1 \in \beta$ and m_2 is a monomial dividing m_1 , then $m_2 \in \beta$. We will call such a set β a **basis set**, and its members **basis monomials**. As a set of $\mathbf{k}$ -points, we define U_β as follows:

$$(4) \quad U_\beta = \{t \in \mathbf{H}^n \mid \mathbf{k}[\mathbf{x}]/I_t \text{ has } \mathbf{k}\text{-basis } \beta\} \subseteq \mathbf{H}^n.$$

Every point $t \in \mathbf{H}^n$ belongs to at least one set of the form U_β , a fact that is, in M. Haiman's words, "often regarded as a part of Gröbner basis theory but actually goes back to Gordan [3]" [6, p. 207]. Therefore, the sets U_β form an open covering of $\mathbf{H}^n$. In fact, the U_β form an open *affine* covering of $\mathbf{H}^n$; this is proved in [9], where the coordinate ring of U_β is explicitly obtained as a quotient of a polynomial ring. Note that Haiman introduced the open subschemes U_β in [6] for the case of the affine plane ($r = 2$); he showed that they are affine using other means.

Remark 2.2.1. Let Z_β denote the restriction of the universal closed subscheme Z_n to $U_\beta \subseteq \mathbf{H}^n$. Then the direct image of the structure sheaf of Z_β on U_β is free with basis β , because β is everywhere a local basis of the (locally free) direct image. This also follows directly from the construction of (U_β, Z_β) given in [9, Sec. 7.3].

Note that the monomials that do *not* belong to a basis set β generate a monomial ideal

$$I = I_\beta \subseteq \mathbf{k}[\mathbf{x}];$$

it is immediate that the quotient $\mathbf{k}[\mathbf{x}]/I_\beta$ has β as $\mathbf{k}$ -basis, that is, I_β corresponds to a point

$$(5) \quad t = t_\beta \in U_\beta.$$

Conversely, if I is a monomial ideal such that $\mathbf{k}[\mathbf{x}]/I$ has $\mathbf{k}$ -dimension n , then the quotient has as $\mathbf{k}$ -basis the set of monomials not in I , and this is clearly a basis set β of n members.

2.3. The coordinate ring of U_β . Let R denote the coordinate ring of the affine scheme U_β , and let $I_{U_\beta} \subseteq R[\mathbf{x}]$ be the ideal that cuts out the universal closed subscheme $Z_\beta \subseteq U_\beta \times \mathbb{A}_\mathbf{k}^r$. By Remark 2.2.1, we have that the module $R[\mathbf{x}]/I_{U_\beta}$ is R -free with basis β . Therefore, every monomial

$$x_1^{d_1} x_2^{d_2} \dots x_r^{d_r} = \mathbf{x}^{\mathbf{d}} \in R[\mathbf{x}]$$

is congruent modulo I_{U_β} to a unique R -linear combination of the monomials $\mathbf{x}^{\mathbf{j}} \in \beta$, or, in other words, for every $\mathbf{x}^{\mathbf{d}}$ there is a unique polynomial of the form

$$(6) \quad F_{\mathbf{d}} = \mathbf{x}^{\mathbf{d}} - \sum_{\mathbf{j} \in \beta} c_{\mathbf{j}}^{\mathbf{d}} \cdot \mathbf{x}^{\mathbf{j}} \in I_{U_\beta}, \quad c_{\mathbf{j}}^{\mathbf{d}} \in R,$$

where we abuse the notation (here and elsewhere) by writing $\mathbf{j} \in \beta$ for $\mathbf{x}^{\mathbf{j}} \in \beta$. When writing monomials in this way, we will reserve $\mathbf{j}$ for basis monomials. Note that

$$(7) \quad \mathbf{d} \in \beta \Rightarrow \begin{cases} c_{\mathbf{j}}^{\mathbf{d}} = 1 \text{ if } \mathbf{d} = \mathbf{j}, \text{ and} \\ c_{\mathbf{j}}^{\mathbf{d}} = 0 \text{ if } \mathbf{d} \neq \mathbf{j}. \end{cases}$$

Following Haiman [6, p. 210], we multiply the polynomial (6) by the variable x_i , and then expand each monomial $x_i \cdot \mathbf{x}^{\mathbf{j}}$ using (6) to obtain a polynomial of the form

$$x_i \cdot \mathbf{x}^{\mathbf{d}} + (R\text{-linear combination of basis monomials}) \in I_{U_\beta},$$

which must therefore be equal to $F_{\mathbf{d}'}$, where $\mathbf{x}^{\mathbf{d}'} = x_i \cdot \mathbf{x}^{\mathbf{d}}$. Equating coefficients, we obtain the relations

$$(8) \quad c_{\mathbf{j}_0}^{\mathbf{d}'} - \sum_{\mathbf{j} \in \beta} c_{\mathbf{j}}^{\mathbf{d}} \cdot c_{\mathbf{j}_0}^{\mathbf{j}'} = 0,$$

where $\mathbf{x}^{\mathbf{j}'} = x_i \cdot \mathbf{x}^{\mathbf{j}}$.

For each coefficient $c_{\mathbf{j}}^{\mathbf{d}}$ such that $\mathbf{d} \notin \beta$ we introduce an indeterminate $C_{\mathbf{j}}^{\mathbf{d}}$, and let

$$(9) \quad \delta: \mathbf{k}[(C_{\mathbf{j}}^{\mathbf{d}})] \rightarrow R, \quad C_{\mathbf{j}}^{\mathbf{d}} \mapsto c_{\mathbf{j}}^{\mathbf{d}},$$

be the natural map. We have the following

Proposition 2.3.1. *The coordinate ring R of U_β is generated as a $\mathbf{k}$ -algebra by the coefficients $c_{\mathbf{j}}^{\mathbf{d}}$ such that $\mathbf{d} \notin \beta$; that is, the map δ is surjective. Furthermore, the kernel of δ is generated by the polynomials*

$$\tau_{\mathbf{j}_0}^{(\mathbf{d}, x_i)} = C_{\mathbf{j}_0}^{\mathbf{d}'} - \sum_{\mathbf{j} \in \beta} C_{\mathbf{j}}^{\mathbf{d}} \cdot C_{\mathbf{j}_0}^{\mathbf{j}'}, \quad \mathbf{d} \notin \beta, \mathbf{j}_0 \in \beta, 1 \leq i \leq r,$$

that are obtained from (8) by replacing each coefficient $c_{\mathbf{j}}^{\mathbf{b}}$ with the corresponding indeterminate $C_{\mathbf{j}}^{\mathbf{b}}$, if $\mathbf{b} \notin \beta$, or the appropriate constant — 0 or 1, according to (7), and denoted $C_{\mathbf{j}}^{\mathbf{b}}$ — if $\mathbf{b} \in \beta$.

Proof: The construction of R given in [9, Sec. 7] shows that R is generated by a certain finite subset of the coefficients $c_{\mathbf{j}}^{\mathbf{d}}$, and the kernel of the map δ is generated by a finite set of polynomials (there denoted $\rho_{\mathbf{j}}^{(\mathbf{b}_1, \mathbf{b}_2)}$), each of which is in fact either equal to one of the polynomials τ or to a $\mathbf{k}$ -linear combination of two of them. $\square$

Remark 2.3.2. It is clear that the point t_β (5) is the “origin” of U_β ; that is, t_β is the point at which all the coefficient functions $c_{\mathbf{j}}^{\mathbf{d}}$ with $\mathbf{d} \notin \beta$ vanish.

2.4. The locus of reduced subschemes meets U_β . Let $\mathbf{H}_o \subseteq \mathbf{H}^n$ denote the open subscheme parameterizing the closed subschemes of $\mathbb{A}_{\mathbf{k}}^r$ that are supported at n distinct points (and are therefore reduced). Since $\mathbf{H}_o$ is irreducible and of dimension rn , its closure $\bar{\mathbf{H}}_o$ is a component of $\mathbf{H}^n$ of dimension rn . It is well-known that the point t_β corresponding to the monomial ideal $I_\beta \subseteq \mathbf{k}[\mathbf{x}]$ lies in $\bar{\mathbf{H}}_o$; in fact, one can exhibit a one-parameter family of length- n closed subschemes of $\mathbb{A}_{\mathbf{k}}^r$ such that the fiber

over 0 is the subscheme defined by I_β and the general fiber is a reduced subscheme. This family is constructed using Hartshorne's concept of **distraction** [8] as adapted by Geramita *et al.* in [2]. The construction works whenever the ground field is infinite (which is true for us, since $\mathbf{k}$ is assumed algebraically closed) or sufficiently large.

Briefly, the construction goes like this. First, choose an infinite (or sufficiently long) sequence of distinct elements $a_0, a_1, a_2, \dots$ of $\mathbf{k}$. Then, for every minimal generator

$$x_1^{d_1} x_2^{d_2} \dots x_r^{d_r} = \mathbf{x}^{\mathbf{d}}$$

of the monomial ideal I_β , form the homogeneous polynomial

$$f_{\mathbf{d}} = \prod_{i=1}^r \left(\prod_{j=0}^{d_i-1} (x_i - a_j \cdot w) \right) \in \mathbf{k}[w][\mathbf{x}].$$

One then proves that the homogeneous ideal $\mathcal{F}$ generated by the polynomials $f_{\mathbf{d}}$ is the ideal of all functions vanishing on the n (distinct projective) points

$$[1 : a_{e_1} : a_{e_2} : \dots : a_{e_r}] \in \mathbb{P}_{\mathbf{k}}^r, \quad x_1^{e_1} x_2^{e_2} \dots x_r^{e_r} = \mathbf{x}^{\mathbf{e}} \in \beta$$

([2, Theorem 2.2]). In particular, if we view w as a parameter, then for any nonzero value w_1 of w , the ideal $\mathcal{F}(w_1) \subset \mathbf{k}[\mathbf{x}]$ obtained by replacing the variable w by the constant w_1 is supported at the n distinct points

$$(w_1 \cdot a_{e_1}, w_1 \cdot a_{e_2}, \dots, w_1 \cdot a_{e_r}), \quad \mathbf{x}^{\mathbf{e}} \in \beta$$

(when $w_1 = 0$, clearly $\mathcal{F}(0) = I_\beta$, the original monomial ideal of colength n). One then checks that the quotient $\mathbf{k}[w][\mathbf{x}]/\mathcal{F}$ is $\mathbf{k}[w]$ -free with basis β as a $\mathbf{k}[w]$ -module. (Indeed, from the form of the polynomials $f_{\mathbf{d}}$, it is clear that every minimal generator of the monomial ideal I_β is congruent (mod $\mathcal{F}$) to a $\mathbf{k}[w]$ -linear combination of basis monomials *that divide the minimal generator*. Induction on the degree then shows that every monomial $m \in I_\beta$ is congruent (mod $\mathcal{F}$) to a $\mathbf{k}[w]$ -linear combination of basis monomials that divide m ; therefore, the basis monomials span the quotient $\mathbf{k}[w][\mathbf{x}]/\mathcal{F}$. Since the $\mathbf{k}$ -dimension of each quotient $\mathbf{k}[\mathbf{x}]/\mathcal{F}(w_1)$ is $\geq n$, we find that β is locally a basis everywhere, and hence a global basis, as desired.) In this way we obtain a finite and flat family of subschemes which defines a map $\varphi: \text{Spec}(\mathbf{k}[t]) \rightarrow U_\beta \subseteq \mathbf{H}^n$; one sees easily that $\varphi(0) = t_\beta$ and $\varphi(w_1) \in \mathbf{H}_\circ \cap U_\beta$ for all $w_1 \neq 0$. It follows at once that $t_\beta \in \bar{\mathbf{H}}_\circ$, which yields

Proposition 2.4.1. *The $\mathbf{k}$ -dimension of the (Zariski) tangent and cotangent spaces of the point t_β is $\geq \dim(\bar{\mathbf{H}}_\circ) = rn$, and t_β is a smooth point of $\mathbf{H}^n$ if and only if*

$$\dim_{\mathbf{k}}(\text{cotangent space of } t_\beta) = rn. \quad \square$$

Remark 2.4.2. We will give another proof that the $\mathbf{k}$ -dimension of the cotangent space of t_β is at least rn in Section 4.

Remark 2.4.3. The foregoing demonstrates that each open subscheme U_β meets the locus $\mathbf{H}_\circ$ of reduced subschemes nontrivially. This is shown in a different way in [9, Sec. 2.4].




FIGURE 2. The “arrows” $c_{(0,0)}^{(1,2)}$, $c_{(0,1)}^{(1,1)}$, $c_{(0,2)}^{(1,1)}$, and $c_{(0,1)}^{(2,0)}$. Monomials not in β are shown in boldface.

3. THE COTANGENT SPACE OF THE POINT $t_\beta \in \mathbf{H}^n$

Recall that the cotangent space of the point $t_\beta \in \mathbf{H}^n$ is the $\mathbf{k}$ -vector space $\mathfrak{m}_{t_\beta}/\mathfrak{m}_{t_\beta}^2$, where $\mathfrak{m}_{t_\beta}$ is the maximal ideal of the local ring $\mathcal{O}_{t_\beta}$. In light of Remark 2.3.2, we see that we can identify the cotangent space with the $\mathbf{k}$ -vector space M/M^2 , where M is the maximal ideal generated by the functions $c_j^{\mathbf{d}}$ ($\mathbf{d} \notin \beta$) in the coordinate ring R of U_β . In [6], Haiman observed, in the two-variable case, that congruence (mod M^2) can be visualized as translation-equivalence of arrows defined by the $c_j^{\mathbf{d}}$. We now extend Haiman’s idea to any number r of variables.

3.1. Arrows, translation, and congruence modulo M^2 . As stated by Haiman, with symbols adjusted, we have that [6, p. 210]

[m]odulo M^2 , the terms $-c_j^{\mathbf{d}} \cdot c_{j_0}^{\mathbf{j}'}$ [in] equation (8) reduce to zero for $\mathbf{j}' \notin \beta$ and for $\mathbf{j}' \in \beta$, $\mathbf{j}' \neq \mathbf{j}_0$ [recall (7) — here we are assuming that $\mathbf{d} \notin \beta$, so that $c_j^{\mathbf{d}} \in M$]. The remaining term is $-c_{j_1}^{\mathbf{d}}$, where

$$\mathbf{j}'_1 = \mathbf{j}_0, \text{ that is, } \mathbf{x}^{\mathbf{j}_1} \cdot x_i = \mathbf{x}^{\mathbf{j}_0},$$

or zero if $\mathbf{x}^{\mathbf{j}_0}$ is not divisible by x_i . Thus in M/M^2 we have

$$(10) \quad c_{j_0}^{\mathbf{d}'} = \begin{cases} c_{j_1}^{\mathbf{d}}, & \text{if } x_i \text{ divides } \mathbf{x}^{\mathbf{j}_0}, \text{ and} \\ 0, & \text{otherwise.} \end{cases}$$

It is convenient to depict each $c_j^{\mathbf{d}}$ by an arrow from $\mathbf{d}$ to $\mathbf{j}$ (see Figure 2). Equation (10) says that we may move these arrows [in the x_i -direction, $1 \leq i \leq r$,] without changing their values modulo M^2 , provided we keep the head inside β and the tail outside β . More generally, as long as we keep the tail in the first [orthant] and outside β , we may even move the head across the [hyperplane (x_i -degree = 0)].

When this is possible, the value of the arrow (mod M^2) is zero.

Henceforth when we speak of an “arrow” $c_j^{\mathbf{d}}$, we will assume that $\mathbf{d} \notin \beta$, so that there is a corresponding indeterminate $C_j^{\mathbf{d}}$ (as defined in Section 2.3). We will say that two arrows $c_{j_1}^{\mathbf{b}_1}$ and $c_{j_2}^{\mathbf{b}_2}$ are **translation-equivalent** if the first can be moved to the second by a sequence of discrete steps in the various variable directions such that

the head of the arrow remains inside β and the tail remains outside β . This clearly defines an equivalence relation on the set of arrows, which we denote $c_{j_1}^{b_1} \sim c_{j_2}^{b_2}$. For example, in Figure 2, we see that $c_{(0,1)}^{(1,1)} \sim c_{(0,2)}^{(1,2)}$. Abusing the language and notation, we say c_j^b is **translation-equivalent** to 0, and write $c_j^b \sim 0$, to indicate that the arrow can be translated to a position such that one more step in some direction of decreasing degree would cause the head of the arrow to exit the first orthant, with the tail remaining a monomial outside of β . The arrow $c_{(0,0)}^{(1,2)}$ in Figure 2 provides an example: one further step in the decreasing x_2 -direction would cause the head of this arrow to exit the first quadrant.

From the foregoing, it is clear that

$$(11) \quad \begin{aligned} c_{j_1}^{b_1} \sim c_{j_2}^{b_2} &\Rightarrow c_{j_1}^{b_1} \equiv c_{j_2}^{b_2} \pmod{M^2}, \text{ and} \\ c_j^b \sim 0 &\Rightarrow c_j^b \equiv 0 \pmod{M^2}. \end{aligned}$$

Furthermore, the reasoning in the quoted passage can be adapted to prove

Proposition 3.1.1. *Let $\tau_{j_0}^{(d, x_i)}$ be any one of the polynomial generators of the kernel of the map δ (9). Then*

- (a) *Each term in $\tau_{j_0}^{(d, x_i)}$ is, up to sign, either a single indeterminate C_j^b or a product of two such indeterminates.*
- (b) *The number of linear terms in $\tau_{j_0}^{(d, x_i)}$ is equal to 1 or 2.*
- (c) *If there are two linear terms in $\tau_{j_0}^{(d, x_i)}$, then these terms have the form $C_{j_0}^{d'}$ and $-C_{j_1}^d$, where $\mathbf{x}^{d'} = x_i \cdot \mathbf{x}^d$ and $\mathbf{x}^{j_0} = x_i \cdot \mathbf{x}^{j_1}$; in particular, the signs differ, and the corresponding arrows $c_{j_0}^{d'}$ and $c_{j_1}^d$ are translation-equivalent (i.e., $c_{j_0}^{d'} \sim c_{j_1}^d$).*
- (d) *If there is only one linear term in $\tau_{j_0}^{(d, x_i)}$, then it is the term $C_{j_0}^{d'}$, and the corresponding arrow $c_{j_0}^{d'} \sim 0$. $\square$*

3.2. k-Linearly independent sets of arrows in M/M^2 . Our main goal in this section is to prove the following

Theorem 3.2.1. *Let S be a finite set of arrows c_j^d having d members. Then S has maximal rank (mod M^2) (that is, the k -span of S in M/M^2 has dimension d) if and only if the following conditions hold:*

- (a) $c_j^d \not\sim 0$ for all $c_j^d \in S$.
- (b) $c_{j_1}^{d_1} \not\sim c_{j_2}^{d_2}$ for all $c_{j_1}^{d_1} \neq c_{j_2}^{d_2}$ in S .

Before proceeding with the proof, we make a few preparations. We first extend the notion of translation-equivalence to the set of indeterminates C_j^b in the obvious way:

$$\begin{aligned} C_{j_1}^{b_1} \sim C_{j_2}^{b_2} &\Leftrightarrow c_{j_1}^{b_1} \sim c_{j_2}^{b_2} \\ C_j^b \sim 0 &\Leftrightarrow c_j^b \sim 0. \end{aligned}$$

Then Proposition 3.1.1 immediately yields

Lemma 3.2.2. *Let $\tau_{j_0}^{(\mathbf{d}, x_i)}$ be any one of the polynomial generators of the kernel of the map δ (9). Then*

- (a) *If $\tau_{j_0}^{(\mathbf{d}, x_i)}$ has two linear terms $C_{j_0}^{\mathbf{d}'}$ and $-C_{j_1}^{\mathbf{d}}$, then $C_{j_0}^{\mathbf{d}'} \sim C_{j_1}^{\mathbf{d}}$.*
- (b) *If $\tau_{j_0}^{(\mathbf{d}, x_i)}$ has just one linear term $C_{j_0}^{\mathbf{d}'}$, then $C_{j_0}^{\mathbf{d}'} \sim 0$. $\square$*

Given a polynomial $P \in \mathbf{k}[(C_j^{\mathbf{b}})]$ of degree q , we write

$$P = P^{(0)} + P^{(1)} + \cdots + P^{(q)},$$

where each $P^{(j)}$ is homogeneous of degree j . We also define the map

$$(12) \quad E: \mathbf{k}[(C_j^{\mathbf{b}})] \rightarrow \mathbf{k}[\langle (C_j^{\mathbf{b}}) \rangle], \quad C_j^{\mathbf{b}} \mapsto \langle C_j^{\mathbf{b}} \rangle,$$

where the target is the ring of polynomials in the translation-equivalence classes $\langle C_j^{\mathbf{b}} \rangle$ of the indeterminates $C_j^{\mathbf{b}}$.

Lemma 3.2.3. *Let $a_1, \dots, a_s \in \mathbf{k}$, and $\tau_1, \dots, \tau_s$ be s of the polynomials $\tau_{j_0}^{(\mathbf{d}, x_i)}$. Let N denote the set of indices n such that τ_n has just one linear term $C_{j_n}^{\mathbf{b}_n}$. Then*

$$E\left(\sum_{j=1}^s a_j \cdot \tau_j^{(1)}\right) = \sum_{n \in N} a_n \cdot \langle C_{j_n}^{\mathbf{b}_n} \rangle.$$

Proof: It suffices to observe that for the τ_j containing two linear terms $C_{j_0}^{\mathbf{b}'_j}$ and $-C_{j_1}^{\mathbf{b}_j}$, we have that

$$E(a_j \cdot \tau_j^{(1)}) = a_j \cdot \left(\langle C_{j_0}^{\mathbf{b}'_j} \rangle - \langle C_{j_1}^{\mathbf{b}_j} \rangle \right) = 0,$$

where the last equality follows from Lemma 3.2.2. $\square$

Proof of Theorem 3.2.1: From the implications (11), it is clear that if either of the conditions (a), (b) in the statement of the theorem fails, then the dimension of the $\mathbf{k}$ -span of S in M/M^2 is $< d$; therefore, if the dimension is d , then (a) and (b) must hold. To prove the converse, we argue by contradiction: Suppose that (a) and (b) hold, but that the $\mathbf{k}$ -span of S in M/M^2 has dimension $< d$. Then there exists a nontrivial $\mathbf{k}$ -linear combination of (distinct) elements of S that is congruent to 0 modulo M^2 , say

$$\sum_{i=1}^m \alpha_i \cdot c_{j_i}^{\mathbf{d}_i} \equiv 0 \pmod{M^2},$$

which implies that

$$\sum_{i=1}^m \alpha_i \cdot c_{j_i}^{\mathbf{d}_i} - (\text{elt. of } M^2) = 0 \in R \quad (R = \text{coord. ring of } U_\beta).$$

From the description of R as a quotient of the polynomial ring $\mathbf{k}[(C_j^{\mathbf{d}})]$ provided by Proposition 2.3.1, we see that we have an equation

$$\sum_{i=1}^m \alpha_i \cdot C_{\mathbf{j}_i}^{\mathbf{d}_i} - (\text{terms in } C_j^{\mathbf{d}} \text{ of degree } \geq 2) = \sum_{j=1}^s g_j \cdot \tau_j,$$

where the coefficients $g_j \in \mathbf{k}[(C_j^{\mathbf{d}})]$ and each τ_j is one of the polynomials $\tau_{\mathbf{j}_0}^{(\mathbf{d}, x_i)}$ that generate the kernel of the map δ (9). Since the τ_j have only linear and quadratic terms (by Proposition 3.1.1), we see that the terms of the form $g_j^{(i)} \cdot \tau_j$ for $i \geq 1$ can be transposed to the left to yield an equation

$$\sum_{i=1}^m \alpha_i \cdot C_{\mathbf{j}_i}^{\mathbf{d}_i} - (\text{terms in } C_j^{\mathbf{d}} \text{ of degree } \geq 2) = \sum_{j=1}^s g_j^{(0)} \cdot \tau_j.$$

Equating the degree-1 terms on both sides, we obtain

$$\sum_{i=1}^m \alpha_i \cdot C_{\mathbf{j}_i}^{\mathbf{d}_i} = \sum_{j=1}^s g_j^{(0)} \cdot \tau_j^{(1)}.$$

As in Lemma 3.2.3, we let N denote the set of indices n such that $\tau_n^{(1)}$ consists of a single term $C_{\mathbf{j}_n}^{\mathbf{b}_n}$. Applying the map E (12) to both sides and rewriting the RHS using the lemma, we find that

$$\sum_{i=1}^m \alpha_i \cdot \langle C_{\mathbf{j}_i}^{\mathbf{d}_i} \rangle = \sum_{n \in N} g_n^{(0)} \cdot \langle C_{\mathbf{j}_n}^{\mathbf{b}_n} \rangle.$$

Recall that conditions (a) and (b) hold, by hypothesis, and that we began with a nontrivial linear combination of the $c_j^{\mathbf{d}} \in S$, so that at least one of the coefficients $\alpha_i \neq 0$. If the corresponding equivalence class $\langle C_{\mathbf{j}_i}^{\mathbf{d}_i} \rangle$ does not appear on the RHS of the last equation, then the term $\alpha_i \cdot \langle C_{\mathbf{j}_i}^{\mathbf{d}_i} \rangle$ must cancel with one or more other terms $\alpha_j \cdot \langle C_{\mathbf{j}_j}^{\mathbf{d}_j} \rangle$ on the LHS; whence, $c_{\mathbf{j}_i}^{\mathbf{d}_i} \sim c_{\mathbf{j}_j}^{\mathbf{d}_j}$, which contradicts condition (b). Therefore, we must have that $\langle C_{\mathbf{j}_i}^{\mathbf{d}_i} \rangle = \langle C_{\mathbf{j}_n}^{\mathbf{d}_n} \rangle$ for some $n \in N$. That is, we have $C_{\mathbf{j}_i}^{\mathbf{d}_i} \sim C_{\mathbf{j}_n}^{\mathbf{b}_n}$, but since $C_{\mathbf{j}_n}^{\mathbf{b}_n} \sim 0$ by Lemma 3.2.2, it follows immediately that $C_{\mathbf{j}_i}^{\mathbf{d}_i} \sim 0 \Leftrightarrow c_{\mathbf{j}_i}^{\mathbf{d}_i} \sim 0$, which contradicts condition (a), and the proof is complete. $\square$

As a corollary, we obtain the following converses to the implications (11):

Corollary 3.2.4. *For $c_j^{\mathbf{b}}$, $c_{\mathbf{j}_1}^{\mathbf{b}_1}$, $c_{\mathbf{j}_2}^{\mathbf{b}_2}$ any arrows, we have that*

- (a) $c_j^{\mathbf{b}} \equiv 0 \pmod{M^2} \Rightarrow c_j^{\mathbf{b}} \sim 0$.
- (b) $c_{\mathbf{j}_1}^{\mathbf{b}_1} \equiv c_{\mathbf{j}_2}^{\mathbf{b}_2} \not\equiv 0 \pmod{M^2} \Rightarrow c_{\mathbf{j}_1}^{\mathbf{b}_1} \sim c_{\mathbf{j}_2}^{\mathbf{b}_2}$.

Proof: Apply the theorem to the sets $\{c_j^{\mathbf{b}}\}$ and $\{c_{\mathbf{j}_1}^{\mathbf{b}_1}, c_{\mathbf{j}_2}^{\mathbf{b}_2}\}$, neither of which has maximal rank $(\text{mod } M^2)$, provided that $c_{\mathbf{j}_1}^{\mathbf{b}_1} \neq c_{\mathbf{j}_2}^{\mathbf{b}_2}$. $\square$

4. A LINEARLY INDEPENDENT SET IN M/M^2 OF CARDINALITY rn

Given a basis set of monomials β of size n in r variables (or equivalently the associated monomial ideal I_β), we exhibit a set $\mathcal{S}$ of arrows c_j^d of cardinality rn whose $\mathbf{k}$ -span in M/M^2 has dimension rn . This gives a second proof that the $\mathbf{k}$ -dimension of the cotangent space of the point $t_\beta \in \mathbf{H}^n$ is at least rn , as promised in Remark 2.4.2. More importantly, if $t_\beta \in \mathbf{H}^n$ is nonsingular, then $\mathcal{S}$ must be a basis of the cotangent space. Because of the particular form of the arrows in $\mathcal{S}$, it is often easy to show in particular cases that $\mathcal{S}$ does *not* span the cotangent space; we conclude that t_β is a singular point in such cases. On the other hand, there are several families of basis sets β for which we can prove that $\mathcal{S}$ is a basis of the cotangent space.

4.1. Standard and minimal arrows. We begin by identifying the type of arrow that will belong to our set $\mathcal{S}$. First we define the **vector** of the arrow c_j^d to be the tuple $\mathbf{j} - \mathbf{d}$ (recall that we only speak of an “arrow” when $\mathbf{d} \notin \beta \Leftrightarrow \mathbf{x}^d \in I_\beta$). Since $\mathbf{x}^d \mid \mathbf{x}^j$ is then impossible, the following is immediate:

Lemma 4.1.1. *For every arrow c_j^d , the vector of the arrow has at least one negative component. $\square$*

We say that c_j^d is a **standard arrow** provided that the vector of the arrow has exactly one negative component; if this component is the i -th, corresponding to the variable x_i , then we say that c_j^d is **standard for x_i** , or **x_i -standard**. Concretely, this means that the monomial $\mathbf{x}^d$ at the tail of the arrow has a strictly larger x_i -degree than the monomial $\mathbf{x}^j$ at the head, but the x_k -degree of $\mathbf{x}^d$ is $\leq$ the x_k -degree of $\mathbf{x}^j$ for all other variables $x_k \neq x_i$. For example, in Figure 2, the arrows $c_{(0,1)}^{(1,1)}$, $c_{(0,2)}^{(1,1)}$, and $c_{(0,1)}^{(2,0)}$ are standard for x_1 , and $c_{(0,0)}^{(1,2)}$ is not a standard arrow.

Recall that the monomials in $x_1, \dots, x_r$ are partially ordered by divisibility, and I_β is generated by the minimal monomials in I_β under this ordering (the **minimal generators** of I_β). We call an arrow c_j^d whose tail $\mathbf{x}^d$ is a minimal generator a **minimal arrow**; if the arrow is standard (for x_i), we call it a **minimal standard arrow (for x_i)**. In light of the results of Section 3, the next lemma shows that in seeking a set of arrows to span the $\mathbf{k}$ -vector space M/M^2 , it suffices to restrict one’s attention to minimal arrows.

Lemma 4.1.2. *Let c_j^d be an arbitrary arrow, and $\mathbf{x}^b$ a minimal generator of I_β such that $\mathbf{x}^b \mid \mathbf{x}^d$ (at least one such minimal generator must clearly exist). Then either $c_j^d \sim 0$ or $c_j^d \sim c_{j_1}^b$ for some $j_1 \in \beta$.*

Proof: We can clearly translate $\mathbf{x}^d$ to $\mathbf{x}^b$ by a series of degree-reducing steps in the various variable directions (each step involves dividing the head and tail of the arrow by one of the x_k). Since dividing a basis monomial (at the head of the arrow) by x_k either keeps us inside β or causes us to exit the first orthant across the hyperplane (x_k -degree = 0), the result follows immediately. $\square$

In fact, the set $\mathcal{S}$ that we are out to construct will consist of minimal *standard* arrows, as our main theorem asserts:

Theorem 4.1.3. *Let β be a basis set of n monomials in the variables $x_1, \dots, x_r$, and $I_\beta \subseteq \mathbf{k}[\mathbf{x}]$ the associated monomial ideal. Then there exists a set $\mathcal{S}$ of minimal standard arrows that has cardinality rn and maximal rank (mod M/M^2); that is, the $\mathbf{k}$ -span of $\mathcal{S}$ in M/M^2 has dimension rn .*

The proof will be given in Section 4.5, after the necessary preparations have been made.

4.2. Advancement of minimal standard arrows. We begin with a host of definitions. Let β, I_β , etc., be as above. Note first of all that, since I_β has finite colength, there is for each variable x_i a minimal exponent $w_i > 0$ such that $x_i^{w_i} \in I_\beta$; we will call w_i the x_i -**width** of β . It is then clear that

$$(13) \quad x_i\text{-degree}(\mathbf{x}^{\mathbf{j}}) < w_i \text{ for all } \mathbf{x}^{\mathbf{j}} \in \beta, \ 1 \leq i \leq r.$$

For example, the basis set shown in Figure 2 has x_1 -width $w_1 = 2$ and x_2 -width $w_2 = 3$.

Given $\mathbf{x}^{\mathbf{j}} \in \beta$, we define the x_i -**column** of $\mathbf{x}^{\mathbf{j}}$ to be the set of monomials $\mathbf{x}^{\mathbf{j}}, \mathbf{x}^{\mathbf{j}}/x_i, \mathbf{x}^{\mathbf{j}}/x_i^2, \dots, \mathbf{x}^{\mathbf{j}}/x_i^q$, where $q = x_i\text{-degree}(\mathbf{x}^{\mathbf{j}})$. Clearly these monomials all belong to β . We define the x_i -**shadow** of an arrow $c_{\mathbf{j}}^{\mathbf{d}}$ to be the set of arrows $c_{\mathbf{j}'}^{\mathbf{d}}$ such that $\mathbf{x}^{\mathbf{j}'}$ is in the x_i -column of $\mathbf{x}^{\mathbf{j}}$, and

$$x_i\text{-height}(c_{\mathbf{j}}^{\mathbf{d}}) = x_i\text{-degree}(\mathbf{x}^{\mathbf{d}}).$$

If $c_{\mathbf{j}}^{\mathbf{d}}$ is a standard arrow for x_i , we define its x_i -**offset** to be the $(r-1)$ -tuple v of non-negative integers obtained by deleting the i -th (negative) component of the vector $\mathbf{j} - \mathbf{d}$.

We say that an x_i -standard arrow $c_{\mathbf{j}}^{\mathbf{d}}$ can be **advanced** if either $c_{\mathbf{j}}^{\mathbf{d}} \sim 0$ or $c_{\mathbf{j}}^{\mathbf{d}} \sim c_{\mathbf{j}_1}^{\mathbf{d}_1}$ with the latter arrow having strictly smaller x_i -height. In each case, the idea is that $c_{\mathbf{j}}^{\mathbf{d}}$ can be translated so that its tail moves closer to the hyperplane ($x_i\text{-degree} = 0$), provided that we allow the head to exit the first orthant across this hyperplane in the first case. Note that the head of an x_i -standard arrow can only be translated out of the first orthant across the hyperplane ($x_i\text{-degree} = 0$), since the tail must remain within the first orthant (and outside of β) during translation.

Lemma 4.2.1. *Suppose that $c_{\mathbf{j}}^{\mathbf{d}}$ is a standard arrow for x_i . Then:*

- (a) *All of the arrows in the x_i -shadow of $c_{\mathbf{j}}^{\mathbf{d}}$ are x_i -standard and have equal offsets.*
- (b) *If $c_{\mathbf{j}}^{\mathbf{d}}$ can be advanced, then so can all the arrows in its x_i -shadow.*
- (c) *If $c_{\mathbf{j}}^{\mathbf{d}} \not\sim 0$ can be advanced, then we can advance this arrow until we reach a minimal standard (for x_i) arrow $c_{\mathbf{j}'}^{\mathbf{d}'} \sim c_{\mathbf{j}}^{\mathbf{d}}$ that cannot be advanced.*

Proof: Statement (a) is immediate. Statement (b) follows from the observation that if c_j^d can be translated to $c_{j_1}^b$, then every arrow in the shadow of c_j^d is simultaneously translated “in parallel” either to an arrow in the shadow of $c_{j_1}^b$ or out of the first orthant (across the hyperplane (x_i -degree = 0)). Statement (c) follows easily from Lemma 4.1.2: after advancing c_j^d to $c_{j_1}^{d_1}$ of smaller x_i -height, we translate by degree-reducing steps to a minimal standard arrow $c_{j'}^{d'}$, and repeat as often as necessary. $\square$

4.3. Shadow promotion. The simultaneous translation and advancement of the arrows in a shadow leads to a process of arrow replacement that is the key to the proof of Theorem 4.1.3; we call this process, which we proceed to describe, *shadow promotion*. Suppose that c_j^d is an x_i -standard arrow that can be advanced, which implies by statement (b) of Lemma 4.2.1 that every arrow in the x_i -shadow of c_j^d can be advanced as well. Consider a sequence of translation steps that advances c_j^d , and let $c_{j_1}^{d_1}$ be the first position in the sequence of steps from which it is possible to move one step in the decreasing x_i -direction either to reach an arrow of x_i -height < x_i -height of c_j^d or to move the head of the arrow out of the first orthant; put another way, $c_{j_1}^{d_1}$ is the first position such that

$$\mathbf{x}^{d_1}/x_i \in I_\beta, \text{ and } x_i\text{-degree}(\mathbf{x}^{d_1}) = x_i\text{-degree}(\mathbf{x}^d).$$

Note that

$$(14) \quad x_i\text{-degree}(\mathbf{x}^{j_1}) = x_i\text{-degree}(\mathbf{x}^j).$$

Let $\mathbf{x}^b$ be a minimal generator of I_β that divides $\mathbf{x}^{d_1}/x_i$; it is clear that

$$x_i\text{-degree}(\mathbf{x}^b) < x_i\text{-degree}(\mathbf{x}^{d_1}).$$

We must have that

$$(15) \quad x_i\text{-degree}(\mathbf{x}^b) > x_i\text{-degree}(\mathbf{x}^{j_1}),$$

since otherwise $x_i\text{-degree}(\mathbf{x}^b) \leq x_i\text{-degree}(\mathbf{x}^{j_1})$ holds in addition to

$$(16) \quad x_k\text{-degree}(\mathbf{x}^b) \leq x_k\text{-degree}(\mathbf{x}^{d_1}) \leq x_k\text{-degree}(\mathbf{x}^{j_1}) \text{ for all } k \neq i,$$

where the last inequality holds because c_j^d and its translate $c_{j_1}^{d_1}$ are standard arrows for x_i . In other words, $\mathbf{x}^b \mid \mathbf{x}^{j_1}$, which is a contradiction since $\mathbf{x}^{j_1}$ is a basis monomial and $\mathbf{x}^b$ is not. The contradiction establishes (15); in particular, we have that $x_i\text{-degree}(\mathbf{x}^b) > 0$.

Furthermore, (16) shows that we can translate the arrow $c_{j_1}^{d_1}$ in degree-reducing directions, excluding the x_i -th, to reach an arrow $c_{j_2}^{d_2}$ such that $\mathbf{x}^{d_2}$ and $\mathbf{x}^b$ differ only in x_i -degree. (Because the arrows involved are x_i -standard, the head of the arrow can never leave the first orthant during any of these steps.) We then have that

$$x_i\text{-degree}(\mathbf{x}^{j_2}) = x_i\text{-degree}(\mathbf{x}^{j_1}) < x_i\text{-degree}(\mathbf{x}^b);$$

therefore, the arrow $c_{j_2}^b$ is a minimal standard arrow for x_i having the same offset as $c_{j_2}^{d_2}$, $c_{j_1}^{d_1}$, and c_j^d , and having the same number of arrows in its shadow as $c_{j_1}^{d_1}$ and c_j^d do (recall (14)). We will call the shadow of $c_{j_2}^b$ the **promotion image** of the shadow of the original standard arrow c_j^d ; the promotion image has the same number of arrows as the original shadow, and the arrows in the promotion image are minimal standard arrows for x_i having strictly smaller x_i -height than the original arrows. Indeed, we can view shadow promotion as the replacement of every arrow $c_{j'}^d$ in the shadow of c_j^d with the arrow $c_{j'_2}^b$ in the shadow of $c_{j_2}^b$ such that

$$(17) \quad x_i\text{-degree}(\mathbf{x}^{j'_2}) = x_i\text{-degree}(\mathbf{x}^{j'}).$$

Since the promotion image depends on the path chosen to advance c_j^d , it is not unique in general.

For example, in Figure 2, the arrow $c_{(0,1)}^{(2,0)}$ is a standard arrow for x_1 that can be advanced; its shadow is the singleton set $\{c_{(0,1)}^{(2,0)}\}$. The promotion image of this shadow is easily seen to be $\{c_{(0,2)}^{(1,1)}\}$.

4.4. Iterated shadow promotion. Select one of the variables x_i , and recall that $x_i^{w_i} = \mathbf{x}^d$ is the least power of x_i that belongs to the monomial ideal I_β ; it is clear that $\mathbf{x}^d$ is a minimal generator of the ideal, which we will call the i -th **corner monomial** of β . Note that *every* arrow c_j^d (with tail at the i -th corner monomial) is minimal and standard for x_i , since only the x_i -degree decreases when we move from the tail to the head of the arrow. In this case, the offset of the arrow is the $(r-1)$ -tuple v obtained by deleting the i -th component of $\mathbf{j}$. Let $\mathcal{S}_0(i, v)$ denote the set of all (minimal x_i -standard) arrows having tail $\mathbf{x}^d$ and offset v , and let $c_{j_1}^d$ be the arrow in $\mathcal{S}_0(i, v)$ whose head $\mathbf{x}^{j_1}$ has maximal x_i -degree $q(i, v)$; it follows that $\mathcal{S}_0(i, v)$ is the x_i -shadow of the arrow $c_{j_1}^d$, and the number of arrows in $\mathcal{S}_0(i, v)$ is $q(i, v) + 1$.

By iterating the process of shadow promotion, we can construct a set of $q(i, v) + 1$ minimal x_i -standard arrows that have offset v and cannot be advanced. Indeed, if none of the arrows in $\mathcal{S}_0(i, v)$ can be advanced, then $\mathcal{S}_0(i, v)$ is itself the desired set, which we will denote $\mathcal{S}(i, v)$. Otherwise, one or more of the arrows in $\mathcal{S}_0(i, v)$ can be advanced; in this case we let $c_{j_2}^d \in \mathcal{S}_0(i, v)$ be the advanceable arrow whose head has maximal x_i -degree, so that the set of all advanceable arrows in $\mathcal{S}_0(i, v)$ is the x_i -shadow of $c_{j_2}^d$, by Lemma 4.2.1. We then promote the x_i -shadow of $c_{j_2}^d$, as described in the previous subsection, and denote the (not necessarily unique) promotion image by $P_0(i, v)$; this permits us to form a new set of minimal x_i -standard arrows of offset v :

$$\mathcal{S}_1(i, v) = (\mathcal{S}_0(i, v) \setminus (x_i\text{-shadow of } c_{j_2}^d)) \cup P_0(i, v).$$

Since the number of arrows in the promotion image is equal to the number of arrows in the shadow being promoted, it is clear that the number of arrows in $\mathcal{S}_1(i, v)$ is equal to the number of arrows in $\mathcal{S}_0(i, v)$. Furthermore, the x_i -height of the arrows

in $P_0(i, v)$ is strictly less than the height of the arrows in $\mathcal{S}_0(i, v)$, which is w_i . If none of the arrows in $\mathcal{S}_1(i, v)$ can be advanced, then $\mathcal{S}_1(i, v)$ is the desired set $\mathcal{S}(i, v)$; otherwise, the set of advanceable arrows in $\mathcal{S}_1(i, v)$ is equal to the shadow of an advanceable arrow $c_{j'}^{d'} \in P_0(i, v)$. Replacing this shadow by its promotion image $P_1(i, v)$, we obtain yet another set of minimal x_i -standard arrows of offset v :

$$\mathcal{S}_2(i, v) = \left(\mathcal{S}_1(i, v) \setminus (x_i\text{-shadow of } c_{j'}^{d'}) \right) \cup P_1(i, v).$$

Continuing in this way, we eventually arrive at the desired set $\mathcal{S}(i, v)$ of minimal x_i -standard arrows of offset v , none of which can be advanced. The process must terminate because at each stage the x_i -height of the arrows that can be advanced is strictly less than the corresponding height at the previous stage, and this height cannot decrease to 0.

4.5. Proof of Theorem 4.1.3. For each of the variables x_i , $1 \leq i \leq r$, we will construct a set $\mathcal{S}(i)$ consisting of n minimal standard arrows for x_i such that no two of the arrows in $\mathcal{S}(i)$ are translation-equivalent to each other, and none is ~ 0 . But then the same is true of

$$\mathcal{S} = \bigcup_{i=1}^r \mathcal{S}(i),$$

since x_i - and x_j -standard arrows cannot be translation-equivalent if $i \neq j$; in particular, the cardinality of $\mathcal{S}$ is rn . Theorem 3.2.1 now yields that the k -span of $\mathcal{S}$ in M/M^2 has dimension rn , as desired.

It remains to construct the sets $\mathcal{S}_i$, but this is not difficult. Simply form the sets $\mathcal{S}(i, v)$ for every possible offset v of a minimal x_i -standard arrow having tail the i -th corner monomial $x_i^{w_i}$, as in Subsection 4.4, and let

$$\mathcal{S}(i) = \bigcup_{\{\text{offsets } v\}} \mathcal{S}(i, v).$$

It is clear that every monomial $\mathbf{x}^j \in \beta$ is the head of an arrow in one of the initial sets $\mathcal{S}_0(i, v)$. Furthermore, the sets $\mathcal{S}(i, v)$ are pairwise disjoint, since arrows of different offsets cannot be equal. Therefore, counting elements in the various sets, we find that

$$\begin{aligned} |\mathcal{S}(i)| &= \sum_{\{\text{offsets } v\}} |\mathcal{S}(i, v)| \\ &= \sum_{\{\text{offsets } v\}} |\mathcal{S}_0(i, v)| \\ &= |\beta| = n. \end{aligned}$$

By construction, the arrows $c_j^b \in \mathcal{S}(i, v)$ are minimal x_i -standard arrows of offset v that cannot be advanced; in particular, we have that $c_j^b \not\sim 0$. Therefore, the set $\mathcal{S}(i)$ consists of n minimal x_i -standard arrows, none of which are ~ 0 . We must now show that no two distinct arrows in $\mathcal{S}(i)$ are translation-equivalent to one another. To this end, let $c_{j_1}^{b_1}$ and $c_{j_2}^{b_2}$ be distinct arrows in $\mathcal{S}(i)$. If these arrows have different offsets, then they cannot possibly be translation-equivalent, so suppose that they

each have offset v ; that is, $c_{j_1}^{b_1}, c_{j_2}^{b_2} \in \mathcal{S}(i, v)$. If the two arrows have different heights (*i.e.*, different tails), then, since neither arrow can be advanced, it is again clear that they cannot be translation-equivalent. If the two arrows have the same height, then they have the same tail; therefore, if they were translation-equivalent, they would be equal, which we are assuming is not the case. We conclude that distinct arrows in $\mathcal{S}(i)$ cannot be translation-equivalent, and the proof is complete. $\square$

Remark 4.5.1. It is clear that none of the arrows in the set $\mathcal{S}$ constructed in Theorem 4.1.3 can be advanced.

Remark 4.5.2. Since the promotion image of a shadow is not necessarily unique, neither are the sets $\mathcal{S}(i)$ and $\mathcal{S}$.

4.6. Summary and terminology. We will call a set $\mathcal{S}(i)$ constructed as in Subsection 4.5 a **standard x_i -sub-bunch**, and the union

$$\mathcal{S} = \bigcup_{i=1}^r \mathcal{S}(i),$$

a **standard bunch**, of arrows for β . These sets have the following properties:

- $\mathcal{S}(i)$: contains $n = |\beta|$ minimal x_i -standard arrows; has maximal rank (mod M^2); and no arrow in the set can be advanced.
- $\mathcal{S}$: is the union of sets $\mathcal{S}(i)$, $1 \leq i \leq r$, and accordingly: contains rn minimal standard arrows; has maximal rank (mod M^2); and no arrow in the set can be advanced.

We will have occasion to consider more general sets of arrows $\mathcal{S}'(i)$ that we call **near-standard x_i -sub-bunches**: these are similar to standard x_i -sub-bunches in that they have cardinality n , have maximal rank (mod M^2), and consist of x_i -standard arrows that cannot be advanced; they differ in that the arrows they contain need not be *minimal*. For example, one way to obtain a near-standard x_i -sub-bunch is to replace some or all of the arrows $c_j^d \in \mathcal{S}(i)$ (a standard x_i -sub-bunch) with arrows

$$c_{j'}^{d'} \sim c_j^d, \text{ such that } x_i\text{-height}(c_{j'}^{d'}) = x_i\text{-height}(c_j^d).$$

We will call the union

$$\mathcal{S}' = \bigcup_{i=1}^r \mathcal{S}'(i),$$

of near-standard sub-bunches $\mathcal{S}'(i)$ a **near-standard bunch** of arrows for β ; it is clear that $\mathcal{S}'$ consists of rn standard arrows that cannot be advanced, and has maximal rank (mod M^2).

We have the following useful corollaries of the proof of Theorem 4.1.3:

Corollary 4.6.1. *Let $c_j^{\mathbf{d}}$ be an x_i -standard arrow for the basis set β . Then every standard x_i -sub-bunch $\mathcal{S}(i)$ of arrows for β contains a unique arrow $c_j^{\mathbf{d}'}$ such that*

$$x_i\text{-degree}(\mathbf{x}^{\mathbf{j}'}) = x_i\text{-degree}(\mathbf{x}^{\mathbf{j}}) \quad \text{and} \quad \text{offset}(c_j^{\mathbf{d}'}) = \text{offset}(c_j^{\mathbf{d}}).$$

Proof: Let

$$p = x_i\text{-degree}(\mathbf{x}^{\mathbf{d}}), \quad m = \mathbf{x}^{\mathbf{d}}/x_i^p, \quad \text{and} \quad \mathbf{x}^{\mathbf{j}_0} = \mathbf{x}^{\mathbf{j}}/m \in \beta.$$

Then $\mathbf{x}^{\mathbf{j}_0}$ has offset v from the corner monomial $x_i^{w_i} = \mathbf{x}^{\mathbf{b}}$, so $c_{j_0}^{\mathbf{b}} \in \mathcal{S}_0(i, v)$, and is the only such arrow whose head has x_i -degree $= x_i\text{-degree}(\mathbf{x}^{\mathbf{j}})$. Under iterated shadow promotion, arrows are replaced by other arrows having the same offset and x_i -degree of the head (17), so we see that any standard x_i -sub-bunch $\mathcal{S}(i)$ contains a unique arrow $c_{j_1}^{\mathbf{d}_1}$ as stated in the lemma, the arrow that ultimately replaces $c_{j_0}^{\mathbf{b}}$. $\square$

Corollary 4.6.2. *If $c_j^{\mathbf{d}}$ is an x_i -standard arrow that cannot be advanced, and whose tail is the corner monomial $\mathbf{x}^{\mathbf{d}} = x_i^{w_i}$, then $c_j^{\mathbf{d}}$ belongs to every standard x_i -sub-bunch $\mathcal{S}(i)$.*

Proof: Clear from the construction. $\square$

5. CONSEQUENCES OF THEOREM 4.1.3

5.1. Necessary and sufficient conditions for $t_\beta \in \mathbf{H}^n$ to be nonsingular. By Proposition 2.4.1, we know that the point $t_\beta \in \mathbf{H}^n$ is nonsingular if and only if the cotangent space M/M^2 has $\mathbf{k}$ -dimension rn . Therefore, if $\mathcal{S}'$ is any standard or near-standard bunch of arrows for β (see Subsection 4.6 for terminology), we have that

$$t_\beta \text{ is nonsingular} \Leftrightarrow \mathcal{S}' \text{ spans } M/M^2 \Leftrightarrow \mathcal{S}' \text{ is a } \mathbf{k}\text{-basis of } M/M^2.$$

We will call β a **smooth** basis set whenever any of these equivalent conditions holds.

Theorem 5.1.1. *In order that the basis set β be smooth, it is necessary and sufficient that the following conditions hold:*

- (a) *Every non-standard arrow $c_j^{\mathbf{b}}$ is translation-equivalent to 0.*
- (b) *There exists a near-standard bunch of arrows $\mathcal{S}'$ such that for any standard arrow $c_j^{\mathbf{d}} \not\sim 0$, there exists an arrow $c_j^{\mathbf{d}'} \in \mathcal{S}'$ such that $c_j^{\mathbf{d}} \sim c_j^{\mathbf{d}'}$.*

Proof: We first prove the necessity. Let $\mathcal{S}' = \mathcal{S}$ be the standard bunch given by Theorem 4.1.3; since β is assumed smooth, $\mathcal{S}'$ is a $\mathbf{k}$ -basis of M/M^2 . Therefore, given a non-standard arrow $c_j^{\mathbf{b}}$, we have that the $\mathbf{k}$ -span in M/M^2 of the enlarged set $\mathcal{S}' \cup \{c_j^{\mathbf{b}}\}$ of $rn + 1$ elements has dimension rn . Since no two distinct arrows in $\mathcal{S}'$ are translation-equivalent, none is translation-equivalent to 0, and no standard arrow can be translation-equivalent to a non-standard arrow, Theorem 3.2.1 yields that $c_j^{\mathbf{b}} \sim 0$. Now let $c_j^{\mathbf{d}} \not\sim 0$ be a standard arrow. If $c_j^{\mathbf{d}} \notin \mathcal{S}'$, then again we must have that the $\mathbf{k}$ -span in M/M^2 of the enlarged set $\mathcal{S}' \cup \{c_j^{\mathbf{d}}\}$ has dimension $< rn + 1$; Theorem 3.2.1 now yields that $c_j^{\mathbf{d}} \sim c_j^{\mathbf{d}'}$ for some $c_j^{\mathbf{d}'} \in \mathcal{S}'$.

It remains to prove the sufficiency; that is, assuming that conditions (a) and (b) hold, we must show that β is smooth, for which it is enough to prove that $\mathcal{S}'$ spans the $\mathbf{k}$ -vector space M/M^2 . However, condition (a) implies that M/M^2 is spanned by the standard arrows, and condition (b) ensures that every standard arrow $c_j^{\mathbf{d}} \not\sim 0$ is in the $\mathbf{k}$ -linear span of $\mathcal{S}'$, so we are done. $\square$

5.2. A sufficient condition for $t_\beta \in \mathbf{H}^n$ to be singular. It is often easy to identify basis sets β for which the conditions (a) and (b) of Theorem 5.1.1 do *not* hold, so that the corresponding point $t_\beta \in \mathbf{H}^n$ is singular; we now present one simple way to do this. (Subsequent sections of the paper will be devoted to identifying basis sets β for which conditions (a) and (b) *do* hold.)

If $\mathbf{x}^j \in \beta$ is maximal among monomials in β for the divisibility ordering, we call $\mathbf{x}^j$ a **maximal basis monomial**. If $\mathbf{x}^{\mathbf{d}}$ is a minimal generator of I_β and $\mathbf{x}^j$ is a maximal basis monomial, we call the arrow $c_j^{\mathbf{d}}$ **rigid**, because we have

Lemma 5.2.1. *An arrow $c_j^{\mathbf{d}}$ such that $\mathbf{x}^{\mathbf{d}}$ is a minimal generator of I_β and $\mathbf{x}^j$ is a maximal basis monomial cannot be translated except to itself. Such an arrow must belong to any $\mathbf{k}$ -basis $\mathcal{B}$ of M/M^2 consisting of arrows.*

Proof: Since the tail of the arrow is a minimal generator of I_β , it is clear that the first step in any nontrivial translation of $c_j^{\mathbf{d}}$ must be in a degree-increasing direction. However, such a step would cause the head of the arrow to exit β and enter I_β , which is forbidden. Therefore, $c_j^{\mathbf{d}}$ cannot be translated except to itself.

Now suppose that $\mathcal{B}$ is a $\mathbf{k}$ -basis of M/M^2 consisting of arrows, and that $c_j^{\mathbf{d}} \notin \mathcal{B}$. Then the $\mathbf{k}$ -span in M/M^2 of the set of arrows $\mathcal{B} \cup \{c_j^{\mathbf{d}}\}$ has dimension $< |\mathcal{B}| + 1$; moreover, $c_j^{\mathbf{d}} \not\sim 0$, since it has no nontrivial translations. It then follows from Theorem 3.2.1 that $c_j^{\mathbf{d}} \sim c_{j_1}^{\mathbf{d}_1}$ for some $c_{j_1}^{\mathbf{d}_1} \in \mathcal{B}$, but then $c_j^{\mathbf{d}} = c_{j_1}^{\mathbf{d}_1}$ by rigidity, which contradicts $c_j^{\mathbf{d}} \notin \mathcal{B}$. We conclude that $c_j^{\mathbf{d}} \in \mathcal{B}$. $\square$

Corollary 5.2.2. *If the basis set β has a non-standard rigid arrow $c_j^{\mathbf{d}}$, then condition (a) of Theorem 5.1.1 is false, so β is not smooth.*

Proof: It is clear that $c_j^{\mathbf{d}}$ is a non-standard arrow that is not translation-equivalent to 0. $\square$

5.3. Example: $\beta = \{1, x_1, x_2, x_3\}$. Figure 3 illustrates the basis set $\beta = \{1, x_1, x_2, x_3\}$, the smallest basis set in three variables that has non-standard rigid arrows, and therefore fails to be smooth by Corollary 5.2.2.

Going further, we recall that by Lemma 4.1.2, the $\mathbf{k}$ -vector space M/M^2 is spanned by minimal arrows. However, in this case, there are $6 \cdot 4 = 24$ minimal arrows, 6 of which are ~ 0 (the ones with head $\mathbf{x}^{(0,0,0)} = 1$), and the other 18 of which are rigid, and so must belong to any basis of M/M^2 consisting of arrows, by Lemma 5.2.1. We conclude from this that the $\mathbf{k}$ -dimension of the cotangent space of t_β is 18. It is




FIGURE 3. The basis set $\beta = \{1, x_1, x_2, x_3\}$, with the minimal generators of I_β shown in boldface. The arrows $c_{(0,1,0)}^{(1,0,1)}$, $c_{(1,0,0)}^{(0,1,1)}$, and $c_{(0,0,1)}^{(1,1,0)}$ are non-standard and rigid, demonstrating that $t_\beta \in \mathbf{H}^4$ is singular.

known (see, *e.g.*, [10, Sec. 1, p. 147], or [11, Sec. 5.1, p. 443]) that $\mathbf{H}^4$ is irreducible of dimension $3 \cdot 4 = 12$, so we reconfirm that β is not smooth.

Finally, note that three of the 18 rigid minimal arrows are non-standard (see Figure 3), leaving 15 rigid minimal standard arrows. It is clear that any near-standard bunch $\mathcal{S}'$ (which contains $3 \cdot 4 = 12$ standard arrows) must exclude at least three of these rigid standard arrows, showing that β also fails to satisfy condition (b) of Theorem 5.1.1.

6. THICKENING OF BASIS SETS

In this section of the paper, we begin to identify certain families of smooth basis sets β . Recall that a basis set is smooth if it is associated to a nonsingular point $t_\beta \in \mathbf{H}^n$, and therefore satisfies conditions (a) and (b) of Theorem 5.1.1. We first consider a natural way to obtain a smooth basis set in r variables by “thickening” a smooth basis set in $r - 1$ variables.

6.1. Definition of thickening. Suppose that β_0 is a basis set of n_0 monomials in the variables $x_1, x_2, \dots, x_{r-1}$, and w_r is a positive integer. We define a basis set in r variables as follows:

$$(18) \quad \beta = \{\mathbf{x}^{\mathbf{j}_0} \cdot x_r^s \mid \mathbf{x}^{\mathbf{j}_0} \in \beta_0, 0 \leq s \leq w_r - 1\};$$

it is indeed easy to check that β is a basis set. We say that β is a **thickening** of β_0 from $r - 1$ to r variables. Note that the number of monomials in β is

$$(19) \quad |\beta| = n = n_0 \cdot w_r.$$

6.2. Minimal generators of I_β .

Lemma 6.2.1. *Let β_0 and β be as above. Then a minimal generator of the ideal I_β is either a minimal generator $\mathbf{x}^{\mathbf{b}}$ of the ideal I_{β_0} (and therefore not divisible by x_r) or else is the corner monomial $x_r^{w_r}$. Furthermore, every arrow with tail $x_r^{w_r}$ is a minimal standard arrow for x_r , and none of these arrows can be advanced.*

Proof: Let $m = \mathbf{x}^{\mathbf{d}}$ be a minimal generator of I_β . If m is not divisible by x_r , then for any other variable x_j , $1 \leq j \leq r-1$, we have that either m/x_j is a monomial in β_0 or is undefined; it follows that m is a minimal generator of $I_{\beta_0} \subseteq \mathbf{k}[x_1, \dots, x_{r-1}]$. If m is divisible by x_r , then $m/x_r \in \beta$. Therefore, $m/x_r = \mathbf{x}^{\mathbf{j}^0} \cdot x_r^s$, where $\mathbf{x}^{\mathbf{j}^0} \in \beta_0$ and $0 \leq s \leq w_r - 1$. In fact, we must have $s = w_r - 1$, since otherwise $m \in \beta$, which contradicts $m \in I_\beta$. But then $x_r^{w_r} \mid m$, and since it is clear that $x_r^{w_r}$ is a minimal generator of β , we must have $m = x_r^{w_r}$. As in Section 4.4, we have that every arrow having tail the corner monomial $x_r^{w_r}$ is a minimal x_r -standard arrow. None of these arrows can be advanced, since $x_r^{w_r}$ is the only minimal generator that is divisible by x_r , and hence is the only minimal generator that can serve as the tail of an x_r -standard arrow; this completes the proof of the lemma. $\square$

6.3. A standard bunch $\mathcal{S}$ for β . Let β be the thickening (18) of β_0 . For the proof of our main result on thickenings, it is convenient to have available a standard bunch $\mathcal{S}$ for β that is closely related to a standard bunch $\mathcal{S}_0$ for β_0 . We proceed to construct such a set of arrows.

Let π denote the projection map taking monomials in $x_1, \dots, x_r$ to monomials in $x_1, \dots, x_{r-1}$ defined by

$$m \mapsto m/x_r^{x_r\text{-deg}(m)}.$$

If $c_{\mathbf{j}}^{\mathbf{d}}$ is an arrow for β such that $\pi(\mathbf{x}^{\mathbf{d}}) \notin \beta_0$, then it is clear that

$$\pi(c_{\mathbf{j}}^{\mathbf{d}}) = c_{\pi(\mathbf{j})}^{\pi(\mathbf{d})}$$

is an arrow for β_0 that we will call the **projection** of $c_{\mathbf{j}}^{\mathbf{d}}$. Furthermore, if we can translate $c_{\mathbf{j}}^{\mathbf{d}}$ to $c_{\mathbf{j}_1}^{\mathbf{d}_1}$ in such a way that the tails of the arrows in the translation path always project to monomials $\notin \beta_0$, then the projections of the arrows define a translation path from $\pi(c_{\mathbf{j}}^{\mathbf{d}})$ to $\pi(c_{\mathbf{j}_1}^{\mathbf{d}_1})$ that only involves steps in the directions $x_1, \dots, x_{r-1}$. In this case we will say that the translation path **projects** from β to β_0 .

For each $c_{\mathbf{j}_0}^{\mathbf{d}_0}$ such that

$$x_r\text{-degree}(\mathbf{x}^{\mathbf{d}_0}) = 0 \quad \text{and} \quad x_r\text{-degree}(\mathbf{x}^{\mathbf{j}_0}) = 0$$

(that is, $c_{\mathbf{j}_0}^{\mathbf{d}_0}$ is an arrow for β_0), define

$$\text{tower}(c_{\mathbf{j}_0}^{\mathbf{d}_0}) = \{c_{\mathbf{j}'}^{\mathbf{d}_0} \mid \mathbf{x}^{\mathbf{j}'} = \mathbf{x}^{\mathbf{j}_0} \cdot x_r^s, \ 0 \leq s \leq w_r - 1\}.$$

It is then clear that any translation of $c_{\mathbf{j}_0}^{\mathbf{d}_0}$ in directions other than the x_r -th can be applied to the entire tower “in parallel”.

Lemma 6.3.1. *For $x_i \neq x_r$, let $c_{\mathbf{j}}^{\mathbf{d}}$ be an x_i -standard arrow for β that can be advanced. Then the projection*

$$\pi(c_{\mathbf{j}}^{\mathbf{d}}) = c_{\mathbf{j}_0}^{\mathbf{d}_0}$$

is an x_i -standard arrow for β_0 that can be advanced; consequently, all the arrows in $\text{tower}(c_{\mathbf{j}_0}^{\mathbf{d}_0})$ can be advanced in parallel.

Proof: We have that

$$x_r\text{-degree}(\mathbf{x}^{\mathbf{d}}) \leq x_r\text{-degree}(\mathbf{x}^{\mathbf{j}}) < r,$$

so the tail $\mathbf{x}^{\mathbf{d}}$ must be divisible by a minimal generator of I_{β_0} , by Lemma 6.2.1. Therefore, the projection $c_{\mathbf{j}_0}^{\mathbf{d}_0}$ is defined. Since the vector $\mathbf{j}_0 - \mathbf{d}_0$ is the same as the original vector $\mathbf{j} - \mathbf{d}$ except in the x_r -component (which is 0 for the former and ≥ 0 for the latter), it follows easily that $c_{\mathbf{j}_0}^{\mathbf{d}_0}$ is x_i -standard. Furthermore, the translation path that advances $c_{\mathbf{j}}^{\mathbf{d}}$ consists entirely of x_i -standard arrows, so it projects from β to β_0 , implying that the projection $c_{\mathbf{j}_0}^{\mathbf{d}_0}$ can be advanced (as an arrow for β_0). As observed earlier, all the arrows in $\text{tower}(c_{\mathbf{j}_0}^{\mathbf{d}_0})$ then advance in parallel. $\square$

We now select a particular standard bunch of arrows $\mathcal{S}$ for β (Theorem 4.1.3 ensures that at least one such set exists). Recall that $\mathcal{S}$ can be constructed as the union of standard x_i -sub-bunches $\mathcal{S}(i)$, $1 \leq i \leq r$; each $\mathcal{S}(i)$ consists of n minimal standard arrows for x_i that cannot be advanced. By the last statement of Lemma 6.2.1, together with Corollary 4.6.2, we have that $\mathcal{S}(r)$ is the set of all n arrows having tail $x_r^{w_r}$. For any $x_i \neq x_r$, we can construct $\mathcal{S}(i)$ as in Subsection 4.5, taking care to translate all towers in parallel, using Lemma 6.3.1. That is, if $c_{\mathbf{j}}^{\mathbf{d}_0}$ is a *minimal* x_i -standard arrow that can be advanced, (so the tail $\mathbf{x}^{\mathbf{d}_0}$ is a minimal generator of β_0 , by Lemma 6.2.1), then the projection

$$\pi(c_{\mathbf{j}}^{\mathbf{d}_0}) = c_{\mathbf{j}_0}^{\mathbf{d}_0}$$

can be advanced (as an arrow for β_0); whence, all the arrows in $\text{tower}(c_{\mathbf{j}_0}^{\mathbf{d}_0})$ (which contains the original arrow $c_{\mathbf{j}}^{\mathbf{d}_0}$) can be advanced in parallel. With this understanding, we see that the construction (for β) of an x_i -standard sub-bunch $\mathcal{S}(i)$ can be carried out as follows: First, restricting to the basis set β_0 and the variables $x_1, \dots, x_{r-1}$, choose a standard x_i -sub-bunch $\mathcal{S}_0(i)$, as in the proof of Theorem 4.1.3. Then set

$$\mathcal{S}(i) = \bigcup_{c_{\mathbf{j}_0}^{\mathbf{b}} \in \mathcal{S}_0(i)} \text{tower}(c_{\mathbf{j}_0}^{\mathbf{b}}).$$

Hence, by setting

$$\mathcal{S}_0 = \cup_{i=1}^{r-1} \mathcal{S}_0(i), \quad \mathcal{S} = \cup_{i=1}^r \mathcal{S}(i),$$

we obtain a standard bunch $\mathcal{S}$ for β that “lies over” a standard bunch $\mathcal{S}_0$ for β_0 in the sense that every arrow in $\mathcal{S}(i)$ projects to an arrow in $\mathcal{S}_0(i)$ for $1 \leq i \leq r-1$.

6.4. Thickenings of smooth basis sets are smooth.

Theorem 6.4.1. *If β_0 is a smooth basis set in $r-1$ variables, then for each integer $w_r > 0$, the thickening β (18) is a smooth basis set.*

Proof: It suffices to show that β satisfies conditions (a) and (b) of Theorem 5.1.1. First suppose that $c_{\mathbf{j}}^{\mathbf{d}}$ is a non-standard arrow for β . If $x_r^{w_r} \mid \mathbf{x}^{\mathbf{d}}$, then we can translate

the arrow $c_j^{\mathbf{d}}$ in degree-decreasing steps so that its tail approaches $x_r^{w_r}$; since no non-standard arrow can have tail $x_r^{w_r}$, we see that at some point the head of the arrow must exit the first orthant, demonstrating that $c_j^{\mathbf{d}} \sim 0$. On the other hand, if $x_r^{w_r} \not\prec \mathbf{x}^{\mathbf{d}}$, then we can assume by Lemmas 4.1.2 and 6.2.1 that we have translated our arrow to $c_{j'}^{\mathbf{b}}$, with tail a minimal generator $\mathbf{x}^{\mathbf{b}}$ of I_{β_0} (we are done if the head of the arrow exits the first orthant during this translation). Let $c_{j_0}^{\mathbf{b}} = \pi(c_{j'}^{\mathbf{b}})$, and observe that $c_{j_0}^{\mathbf{b}}$ is a non-standard arrow for β_0 . Since β_0 is a smooth basis set, by hypothesis, we have that $c_{j_0}^{\mathbf{b}} \sim 0$, using only translations in the first $r-1$ variable directions. It is clear that the arrow $c_{j'}^{\mathbf{b}}$ translates “in parallel” with $c_{j_0}^{\mathbf{b}}$, leading to the conclusion that $c_{j'}^{\mathbf{b}} \sim 0$. Therefore β satisfies condition (a) of Theorem 5.1.1.

To prove that β satisfies condition (b) of Theorem 5.1.1, we use the standard bunches $\mathcal{S}$ and $\mathcal{S}_0$ constructed in Subsection 6.3, and we let $c_j^{\mathbf{d}} \not\sim 0$ be an x_i -standard arrow for β . If $i = r$, then we can translate in degree-reducing steps until we reach an arrow whose tail is the corner monomial $\mathbf{x}^{\mathbf{d}_1} = x_r^{w_r}$, so that $c_j^{\mathbf{d}} \sim c_{j_1}^{\mathbf{d}_1} \in \mathcal{S}(r) \subseteq \mathcal{S}$. If $i \neq r$, then we can translate by degree-reducing steps until we reach an arrow $c_{j_1}^{\mathbf{b}}$ whose tail is one of the minimal generators $\mathbf{x}^{\mathbf{b}}$ of I_{β_0} . Let $c_{j_0}^{\mathbf{b}} = \pi(c_{j_1}^{\mathbf{b}})$, and note that $c_{j_1}^{\mathbf{b}} \in \text{tower}(c_{j_0}^{\mathbf{b}})$. Since β_0 is a smooth basis set, we know that there exists an arrow $c_{j'_0}^{\mathbf{b}'} \in \mathcal{S}_0(i)$ such that $c_{j_0}^{\mathbf{b}} \sim c_{j'_0}^{\mathbf{b}'}$. The translations only involve the variable directions $x_1, \dots, x_{r-1}$, and will carry $c_{j_1}^{\mathbf{b}}$ “in parallel” to an arrow $c_{j'_1}^{\mathbf{b}'} \in \text{tower}(c_{j'_0}^{\mathbf{b}'}) \subseteq \mathcal{S}(i)$. We conclude that β satisfies condition (b) of Theorem 5.1.1. This completes the proof that the thickening β of β_0 is a smooth basis set. $\square$

Remark 6.4.2. The converse of Theorem 6.4.1 also holds; that is, if β is a smooth basis set that is a thickening (18) of β_0 , then β_0 is smooth. We leave this as an exercise for the reader.

6.5. Example: “Boxes” are smooth basis sets. Let $w_1, \dots, w_r$ be positive integers, and let

$$(20) \quad \mathcal{B}(w_1, \dots, w_r) = \{x_1^{d_1} x_2^{d_2} \dots x_r^{d_r} \mid 0 \leq d_i < w_i, 1 \leq i \leq r\},$$

which is clearly a basis set containing

$$|\mathcal{B}(w_1, \dots, w_r)| = n = \prod_{i=1}^r w_i$$

monomials; for obvious reasons, we call this type of basis set a **box**. The following useful results are immediate:

Lemma 6.5.1. *The minimal generators of $I_{\mathcal{B}(w_1, \dots, w_r)}$ are the corner monomials $x_i^{w_i}$, $1 \leq i \leq r$. $\square$*

Lemma 6.5.2. *If $m_1, m_2 \in \mathcal{B}(w_1, \dots, w_r)$, then*

$$\text{lcm}(m_1, m_2) \in \mathcal{B}(w_1, \dots, w_r). \quad \square$$

We could give an easy proof of the next proposition using Lemma 6.5.1, but instead we offer a proof based on thickenings.

Proposition 6.5.3. *The box $\mathcal{B}(w_1, \dots, w_r)$ is a smooth basis set.*

Proof: It is clear that $\mathcal{B}(w_1, \dots, w_r)$ is a thickening of $\mathcal{B}(w_1, \dots, w_{r-1})$, so if the latter is smooth, then so is the former, by Theorem 6.4.1. The desired result therefore follows by induction, provided that the result holds in the base case: boxes $\beta = \mathcal{B}(w_1)$ in one variable. But in this case, the monomial ideal I_β has just one minimal generator $x_1^{w_1}$; therefore, there are exactly $w_1 = |\beta|$ minimal arrows, all of which are standard and cannot be advanced; in fact, the set of minimal arrows is equal to the standard bunch $\mathcal{S} = \mathcal{S}(1)$, in the notation of the proof of Theorem 4.1.3. Since M/M^2 is spanned by minimal arrows (Lemma 4.1.2), we conclude that $\mathcal{S}$ is a $\mathbf{k}$ -basis of $M/M^2 \Rightarrow \beta$ is smooth, as desired. $\square$

7. TRUNCATION OF BASIS SETS

Truncation is another natural way to obtain a basis set from a given basis set. In this section we will define the truncation operation and establish one of its key properties: If β is such that every non-standard arrow $c_j^{\mathbf{d}}$ is translation-equivalent to 0 (that is, β satisfies condition (a) of Theorem 5.1.1), then any truncation of β also has this property; in particular, this is so if β is smooth. Under an additional hypothesis, one can further prove that certain truncations of a smooth basis set are smooth (that is, also satisfy condition (b) of Theorem 5.1.1); we discuss this in the next section.

7.1. Definition of truncation. Let β be a basis set of n monomials in the variables $x_1, \dots, x_r$. Choose one of the variables x_j , and a positive integer h such that β contains at least one monomial that is divisible by x_j^h . Then we define the x_j -**truncation of β at height h** to be the basis set

$$(21) \quad \beta_t(x_j, h) = \beta_t = \{m \mid x_j^h \cdot m \in \beta\}.$$

That is, β_t is obtained by discarding the monomials in β with x_j -degree $< h$, and dividing the remaining monomials by x_j^h . Figure 4 provides an example. We write

$$n = |\beta|, \quad n_t = |\beta_t|$$

for the number of monomials in β , β_t , respectively.

7.2. Minimal generators of I_{β_t} . We denote by w_i (resp. $w_i^{(t)}$) the x_i -width of β (resp. β_t); that is, $x_i^{w_i}$ (resp. $x_i^{w_i^{(t)}}$) is the i -th corner monomial of I_β (resp. I_{β_t}). We have the following

Lemma 7.2.1. *Let β be a basis set, and β_t the x_j -truncation of β at height h . Then*

x_2^6			
x_2^5	$x_1x_2^5$		
x_2^4	$x_1x_2^4$	$x_1^2x_2^4$	$x_1^3x_2^4$
x_2^3	$x_1x_2^3$	$x_1^2x_2^3$	$x_1^3x_2^3$
x_2^2	$x_1x_2^2$	$x_1^2x_2^2$	$x_1^3x_2^2$
x_2	x_1x_2	$x_1^2x_2$	$x_1^3x_2$
1	x_1	x_1^2	x_1^3

FIGURE 4. The small basis set enclosed by the polygon is the truncation at x_1 -degree 3 of the basis set consisting of all of the displayed monomials. It is obtained by dividing the monomials to the right of the vertical line by x_1^3 .

- (a) If m is a minimal generator of I_β such that $x_j\text{-degree}(m) \geq h$, then $m/x_j^h = m_t$ is a minimal generator of I_{β_t} ; consequently, for all $x_k \neq x_j$, we have that

$$x_k\text{-degree}(m) = x_k\text{-degree}(m_t) \leq w_k^{(t)}.$$

- (b) The minimal generators m of I_β that have $x_j\text{-degree} > h$ are in bijective correspondence with the minimal generators $m_t = m/x_j^h$ of I_{β_t} that are divisible by x_j .
- (c) If m'_t is a minimal generator of I_{β_t} that is not divisible by x_j , then $x_j^h \cdot m'_t = m'$ is an x_j -multiple of a minimal generator of I_β .
- (d) $w_j = w_j^{(t)} + h > h$, and for all $x_k \neq x_j$, $w_k \geq w_k^{(t)}$.

Proof: We begin by noting that a monomial $m \in I_\beta$ (resp. I_{β_t}) is a minimal generator if and only if for each variable x_i , either $x_i \nmid m$ or $m/x_i \in \beta$ (resp. β_t).

Let m and m_t be as in assertion (a). Since $x_j^h \cdot m_t = m \notin \beta$, we have that $m_t \notin \beta_t \Rightarrow m_t \in I_{\beta_t}$. For any variable $x_k \neq x_j$, note that

$$x_k \mid m_t \Rightarrow x_k \mid m \Rightarrow m/x_k \in \beta;$$

moreover,

$$x_j\text{-degree}(m/x_k) = x_j\text{-degree}(m) \geq h;$$

therefore, $m_t/x_k \in \beta_t$. If $x_j \mid m_t$, then

$$x_j\text{-degree}(m) = (x_j\text{-degree}(m_t) + h) > h;$$

whence,

$$x_j\text{-degree}(m/x_j) \geq h, \text{ and as before } m/x_j \in \beta,$$

which yields $m_t/x_j \in \beta_t$. This completes the proof that m_t is a minimal generator of I_{β_t} , and the stated consequence is immediate.

Since the map $m \mapsto m_t$ is injective, to prove assertion (b), it remains to show that if m'_t is a minimal generator of β_t that is divisible by x_j , then $x_j^h \cdot m'_t = m'$ is a minimal

generator of β . It is clear that $m' \in I_\beta$, since $m'_t \notin \beta_t$. Furthermore, for any variable x_i , we have that

$$(22) \quad x_i | m' \Rightarrow x_i | m'_t \Rightarrow m'_t / x_i \in \beta_t \Rightarrow m' / x_i \in \beta$$

(the first implication is trivial for $x_i = x_j$), so m' is indeed a minimal generator of β .

Let now m'_t, m' be as in assertion (c). As in the preceding paragraph, $m' \in I_\beta$. We can translate m' to a minimal generator m of I_β by degree-reducing steps, but since the implications (22) hold for all $x_i \neq x_j$, we see that we can only move in the direction of decreasing x_j -degree; whence, m' is an x_j -multiple of m , as desired.

Turning to assertion (d), it is clear that

$$\begin{aligned} x_\ell | x_j^{w_j^{(t)}+h} &\Rightarrow x_\ell = x_j \\ &\Rightarrow x_j^{w_j^{(t)}+h} / x_\ell = x_j^{w_j^{(t)}+h-1} = x_j^h \cdot x_j^{w_j^{(t)}-1} \in x_j^h \cdot \beta_t \subseteq \beta, \end{aligned}$$

which implies that $x_j^{w_j^{(t)}+h}$ is a minimal generator of I_β ; therefore, $w_j = w_j^{(t)} + h$. Finally, for any $x_k \neq x_j$, we have that

$$\begin{aligned} x_k^{w_k^{(t)}-1} \in \beta_t &\Rightarrow x_j^h \cdot x_k^{w_k^{(t)}-1} \in \beta \\ &\Rightarrow x_k^{w_k} \nmid (x_j^h \cdot x_k^{w_k^{(t)}-1}) \\ &\Rightarrow w_k \geq w_k^{(t)}, \end{aligned}$$

and the proof is complete. $\square$

7.3. Lifting arrows from β_t to β . Let β and β_t be as above, and let $c_j^{\mathbf{d}}$ be an arrow for β_t . Since $\mathbf{x}^{\mathbf{j}} \in \beta_t$ (resp. $\mathbf{x}^{\mathbf{d}} \notin \beta_t$), we have that $\mathbf{x}^{\mathbf{j}'} = \mathbf{x}^{\mathbf{j}} \cdot x_j^h \in \beta$ (resp. $\mathbf{x}^{\mathbf{d}'} = \mathbf{x}^{\mathbf{d}} \cdot x_j^h \notin \beta$); therefore, $c_j^{\mathbf{d}'}$ is an arrow for β . We will call $c_j^{\mathbf{d}'}$ the **lifting** of $c_j^{\mathbf{d}}$ from β_t to β , and say that $c_j^{\mathbf{d}'}$ **descends** to $c_j^{\mathbf{d}}$. We extend this terminology to sets of arrows in the obvious way. It is clear that the arrows $c_j^{\mathbf{d}}$ and $c_j^{\mathbf{d}'}$ have the same vector

$$\mathbf{j} - \mathbf{d} = \mathbf{j}' - \mathbf{d}'.$$

7.4. Non-standard arrows on β and β_t .

Theorem 7.4.1. *Let β be a basis set with the property that every one of its non-standard arrows is translation-equivalent to 0 (condition (a) of Theorem 5.1.1), and let β_t be the x_j -truncation of β at height h (21). Then β_t also has this property; that is, every non-standard arrow for β_t is translation-equivalent to 0.*

Proof: Let $c_j^{\mathbf{d}}$ be a non-standard arrow for β_t . We must show that $c_j^{\mathbf{d}} \sim 0$. To do this, we consider the lifting $c_j^{\mathbf{d}'}$ of $c_j^{\mathbf{d}}$ to β . Suppose first that

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{d}}) \leq x_j\text{-degree}(\mathbf{x}^{\mathbf{j}});$$

that is, the j -th component of the vector $\mathbf{j} - \mathbf{d}$ is ≥ 0 . By hypothesis, we can translate the arrow $c_j^{\mathbf{d}'}$ so that the head eventually leaves the first orthant. In fact, we *claim*

that we can perform this translation without ever passing through a position $c_{j_1}^{\mathbf{d}_1}$ for which

$$x_j\text{-height}(c_{j_1}^{\mathbf{d}_1}) < x_j\text{-height}(c_{j'}^{\mathbf{d}'}) = x_j\text{-degree}(\mathbf{x}^{\mathbf{d}'}).$$

Proof of claim: We argue by induction on the x_j -height of the original arrow $c_{j'}^{\mathbf{d}'}$. The case of height 0 is trivial: no arrows can have x_j -height < 0 . So suppose that

$$x_j\text{-height}(c_{j'}^{\mathbf{d}'}) = p > 0,$$

and that the claim holds for all heights $< p$. Consider a sequence of steps that will translate the arrow's head out of the first orthant. Let $c_{j_0}^{\mathbf{d}_0}$ be the first point on this path (if any) from which translation to an x_j -height $< p$ is possible; that is,

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{d}_0}) = p, \text{ and } \mathbf{x}^{\mathbf{d}_0}/x_j \in I_\beta.$$

Let $s \geq 1$ be the largest integer such that

$$\mathbf{x}^{\mathbf{b}} = \mathbf{x}^{\mathbf{d}_0}/x_j^s \in I_\beta.$$

Then the arrow $c_{j_0}^{\mathbf{b}}$ is non-standard with the same negative and non-negative components in its vector as the original arrow (the only change in the vector is an increased j -th component, which was ≥ 0 to begin with). In light of the hypothesis on β , and our induction hypothesis, we can translate the arrow $c_{j_0}^{\mathbf{b}}$ (of x_j -height $q < p$) so that the head exits the first orthant without ever reaching a position of height $< q$. It is now clear that the arrow $c_{j_0}^{\mathbf{d}_0}$ can be translated “in parallel” with the arrow $c_{j_0}^{\mathbf{b}}$, and the height of the former never becomes $< p$. This shows that the claim holds for any non-standard arrow at x_j -height p , which completes the proof of the claim.

Returning to the proof of Theorem 7.4.1, we now see that the lifted arrow $c_{j'}^{\mathbf{d}'}$ can be translated so that the head leaves the first orthant and no position on the path has x_j -height less than the original height (which is $\geq h$). By descending the path to β_t , we obtain that the original arrow $c_j^{\mathbf{d}}$ is translation-equivalent to 0 for the basis set β_t .

It remains to show that the same conclusion holds when the j -th component of the vector of our non-standard arrow $c_j^{\mathbf{d}}$ is < 0 ; that is,

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{d}}) > x_j\text{-degree}(\mathbf{x}^{\mathbf{j}}).$$

The lifted arrow $c_{j'}^{\mathbf{d}'}$ is by hypothesis translation-equivalent to 0 on β . Note that any step $c_{j_1}^{\mathbf{d}_1}$ on the corresponding translation path such that

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) \geq h$$

descends to β_t . Therefore, by descending the initial segment of the path, up to the first position $c_{j_1}^{\mathbf{d}_1}$ for which the $x_j\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) < h$ (if any), we conclude that $c_j^{\mathbf{d}}$ is translation-equivalent to 0 for β_t , and we are done. $\square$

8. SUFFICIENT CONDITIONS FOR A TRUNCATION TO BE SMOOTH

Let β be a *smooth* basis set in the variables $x_1, \dots, x_r$, and β_t the x_j -truncation of β at height h (21). In the last section (Theorem 7.4.1) we saw that β_t necessarily satisfies condition (a) of Theorem 5.1.1. Our main goal in this section is to prove Theorem 8.6.1, which states that β_t will also satisfy condition (b) of Theorem 5.1.1, and therefore be smooth, if we assume an additional hypothesis. Much of our work involves the construction of a (near-)standard x_i -sub-bunch of arrows for β that contains the lifting of a standard x_i -sub-bunch of arrows for β_t . There are two cases: $x_i = x_j$ (the easier case), discussed in Subsection 8.2, and $x_i = x_k \neq x_j$, discussed in Subsection 8.5.

8.1. The additional hypothesis. By the first assertion of Lemma 7.2.1, we know that if m is any minimal generator of I_β , then

$$\forall x_k \neq x_j, \ x_j\text{-degree}(m) \geq h \Rightarrow x_k\text{-degree}(m) \leq w_k^{(t)}.$$

To prove Theorem 8.6.1, we need to control $x_k\text{-degree}(m)$ when $x_j\text{-degree}(m) < h$, using the following

Hypothesis 8.1.1. *For every minimal generator m of I_β , we have that*

$$\forall x_k \neq x_j, \ x_j\text{-degree}(m) < h \text{ and } x_k \mid m \Rightarrow x_k\text{-degree}(m) \geq w_k^{(t)}.$$

8.2. x_j -sub-bunches of arrows for β and β_t .

Lemma 8.2.1. *Let β be an arbitrary (not necessarily smooth) basis set such that β_t , the x_j -truncation at height h (21), is defined. Let $\mathcal{S}(j)$ be a standard x_j -sub-bunch of arrows for β , constructed as in the proof of Theorem 4.1.3. Then the set of arrows $c_j^{\mathbf{d}} \in \mathcal{S}(j)$ with heads $\mathbf{x}^{\mathbf{j}}$ divisible by x_j^h (so that $\mathbf{x}^{\mathbf{j}}/x_j^h \in \beta_t$) is the lifting of a standard x_j -sub-bunch of arrows $\mathcal{S}^{(t)}(j)$ for β_t .*

Proof: Consider a minimal x_j -standard arrow $c_j^{\mathbf{d}}$ for β such that

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{j}}) \geq h \Rightarrow x_j\text{-degree}(\mathbf{x}^{\mathbf{d}}) > h,$$

and suppose that $c_j^{\mathbf{d}}$ can be advanced. We can then promote the entire x_j -shadow of $c_j^{\mathbf{d}}$, as described in Subsection 4.3; the promotion image is the x_j -shadow of an arrow $c_{j_2}^{\mathbf{b}}$, where $\mathbf{x}^{\mathbf{b}}$ is a minimal generator of I_β ,

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{b}}) > x_j\text{-degree}(\mathbf{x}^{\mathbf{j}_2}) = x_j\text{-degree}(\mathbf{x}^{\mathbf{j}}) \geq h$$

and

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{b}}) < x_j\text{-degree}(\mathbf{x}^{\mathbf{d}}).$$

Let $c_{j_t}^{\mathbf{d}_t}$ (resp. $c_{j_{2,t}}^{\mathbf{b}_t}$) denote the arrow that $c_j^{\mathbf{d}}$ (resp. $c_{j_2}^{\mathbf{b}}$) descends to (on the truncation β_t). Lemma 7.2.1 implies that $\mathbf{x}^{\mathbf{d}_t}$ and $\mathbf{x}^{\mathbf{b}_t}$ are minimal generators of I_{β_t} ; moreover, it is clear that the x_j -shadow of $c_{j_{2,t}}^{\mathbf{b}_t}$ is the promotion image of the shadow of $c_{j_t}^{\mathbf{d}_t}$,

and that the lifting of the shadow of $c_{\mathbf{j}_t}^{\mathbf{d}_t}$ (resp. $c_{\mathbf{j}_{2,t}}^{\mathbf{b}_t}$) to β consists of the arrows in the shadow of $c_{\mathbf{j}}^{\mathbf{d}}$ (resp. $c_{\mathbf{j}_2}^{\mathbf{b}}$) whose heads have x_j -degree $\geq h$. Applying this observation to the iterated shadow promotion operations used to construct $\mathcal{S}(j)$, as described in Subsection 4.4, one sees that the desired result follows readily. $\square$

8.3. x_k -standard arrows of x_j -height $\geq h$. Recall from Subsection 4.4 that $\mathcal{S}_0(k, v)$ denotes the set of all (minimal x_k -standard) arrows for β having tail $x_k^{w_k}$ and offset v . Suppose that $c_{\mathbf{j}}^{\mathbf{d}} \in \mathcal{S}_0(k, v)$ has head $\mathbf{x}^{\mathbf{j}} \in \beta_t$. It is then clear that the entire x_k -shadow of $c_{\mathbf{j}}^{\mathbf{d}}$ has this property; therefore, the subset of all arrows in $\mathcal{S}_0(k, v)$ with heads in β_t is equal to the x_k -shadow of $c_{\mathbf{j}_v}^{\mathbf{d}}$, the member of the subset whose head has maximal x_k -degree

$$(23) \quad r(k, v) = x_k\text{-degree}(\mathbf{x}^{\mathbf{j}_v}).$$

If no arrow in $\mathcal{S}_0(k, v)$ has head in β_t , we define $r(k, v) = -1$, and $\mathbf{x}^{\mathbf{j}_v}$ is undefined. Recalling that the x_j -height of an arrow is the x_j -degree of its tail, we have the following

Lemma 8.3.1. *Let $c_{\mathbf{j}}^{\mathbf{b}}$ be an x_k -standard arrow for β having x_j -height $\geq h$ and offset v . Then*

$$x_k\text{-degree}(\mathbf{x}^{\mathbf{j}}) \leq r(k, v).$$

Proof: If not, then, since $c_{\mathbf{j}}^{\mathbf{b}}$ is x_k -standard, we have that

$$\begin{aligned} x_j\text{-degree}(\mathbf{x}^{\mathbf{j}}) &\geq x_j\text{-degree}(\mathbf{x}^{\mathbf{b}}) \geq h, \text{ and} \\ x_i\text{-degree}(\mathbf{x}^{\mathbf{j}}) &\geq x_i\text{-degree}(\mathbf{x}^{\mathbf{b}}), \quad i \neq j, \quad i \neq k. \end{aligned}$$

Let

$$p = x_k\text{-degree}(\mathbf{x}^{\mathbf{b}}), \quad m = \mathbf{x}^{\mathbf{b}}/x_k^p, \quad \text{and} \quad \mathbf{x}^{\mathbf{j}_1} = \mathbf{x}^{\mathbf{j}}/m \in \beta.$$

Then we have that $\mathbf{x}^{\mathbf{j}_1}$ has offset v from the corner monomial $x_k^{w_k} = \mathbf{x}^{\mathbf{d}}$, and

$$x_k\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) = x_k\text{-degree}(\mathbf{x}^{\mathbf{j}}).$$

We further have that $\mathbf{x}^{\mathbf{j}_1} \in \beta_t$ because

$$x_j^h \mid m \Rightarrow (x_j^h \cdot \mathbf{x}^{\mathbf{j}_1}) \mid (m \cdot \mathbf{x}^{\mathbf{j}_1}), \text{ and } m \cdot \mathbf{x}^{\mathbf{j}_1} = \mathbf{x}^{\mathbf{j}} \in \beta, \text{ so } x_j^h \cdot \mathbf{x}^{\mathbf{j}_1} \in \beta.$$

It then follows from the definition (23) of $r(k, v)$ that

$$x_k\text{-degree}(\mathbf{x}^{\mathbf{j}}) = x_k\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) \leq r(k, v),$$

as desired. $\square$

8.4. Linear independence of lifts of x_k -sub-bunches.

Lemma 8.4.1. *Let β and β_t be as above, and let $x_k \neq x_j$. Suppose in addition that the specialization of Hypothesis 8.1.1 to x_k holds; that is, for every minimal generator m of I_β , we have that*

$$(24) \quad x_j\text{-degree}(m) < h \text{ and } x_k \mid m \Rightarrow x_k\text{-degree}(m) \geq w_k^{(t)}.$$

Then, given two x_k -standard arrows (for β) $c_{j_1}^{d_1} \sim c_{j'_1}^{d'_1}$ of x_j -height $\geq h$, there is a translation path from $c_{j_1}^{d_1}$ to $c_{j'_1}^{d'_1}$ such that every arrow in the path has x_j -height $\geq h$.

Proof: Consider a translation path from $c_{j_1}^{d_1}$ to $c_{j'_1}^{d'_1}$ that at some point involves an arrow of x_j -height $< h$, and let $c_{j_2}^{d_2}$ be the arrow in the path that immediately precedes the very first such arrow; in particular, since the next step must be in the negative x_j -direction to reach x_j -height $h - 1$, we have that

$$\mathbf{x}^{d_2}/x_j \in I_\beta, \text{ and } x_j\text{-degree}(\mathbf{x}^{d_2}/x_j) = h - 1.$$

There exists a minimal generator m of I_β such that

$$m \mid (\mathbf{x}^{d_2}/x_j) \Rightarrow x_j\text{-degree}(m) < h.$$

Furthermore, we have that $x_k \mid m$, since the reasoning leading to the inequality (15) implies that

$$x_k\text{-degree}(m) > x_k\text{-degree}(\mathbf{x}^{j_2}) \geq 0,$$

otherwise we have the contradiction $m \mid \mathbf{x}^{j_2} \Rightarrow m \in \beta$.

The hypothesis (24) now implies that

$$w_k^{(t)} \leq x_k\text{-degree}(m) \leq x_k\text{-degree}(\mathbf{x}^{d_2}/x_j) = x_k\text{-degree}(\mathbf{x}^{d_2}).$$

Indeed, similar reasoning shows that any arrow in the path having x_j -height $< h$ must have x_k -height $\geq w_k^{(t)}$. Since the path terminates in the arrow $c_{j'_1}^{d'_1}$ of x_j -height $\geq h$, the path must eventually reach an arrow $c_{j_3}^{d_3}$ following $c_{j_2}^{d_2}$ such that the x_j -height of $c_{j_3}^{d_3}$ equals h , all subsequent arrows in the path have x_j -height $\geq h$, and the arrow preceding $c_{j_3}^{d_3}$ has x_j -height $= h - 1$, so that the step to $c_{j_3}^{d_3}$ is in the increasing x_j -direction. An argument similar to that for $\mathbf{x}^{d_2}$ yields that $x_k\text{-degree}(\mathbf{x}^{d_3}) \geq w_k^{(t)}$. It follows that both $\mathbf{x}^{d_2}$ and $\mathbf{x}^{d_3}$ are divisible by $\mathbf{x}^b = x_j^h \cdot x_k^{w_k^{(t)}} \in I_\beta$; therefore, each of the arrows $c_{j_2}^{d_2}$ and $c_{j_3}^{d_3}$ can be translated by degree-reducing steps to an arrow $c_{j_4}^b$. The heads of the arrows cannot leave the first orthant during these translations because, the arrows being x_k -standard (and of x_k -height $\geq w_k^{(t)}$), the exit would have to be across the hyperplane ($x_k\text{-degree} = 0$), but then the original arrow $c_{j_1}^{d_1}$ of x_k -height $\leq w_k^{(t)}$ could not have existed. We now replace the original path segment from $c_{j_2}^{d_2}$ to $c_{j_3}^{d_3}$ by the translation from $c_{j_2}^{d_2}$ to $c_{j_4}^b$ and the reversal of the translation from $c_{j_3}^{d_3}$ to $c_{j_4}^b$.

Since the latter translations only involve arrows of x_j -height $= h$, we have produced a path from $c_{j_1}^{\mathbf{d}_1}$ to $c_{j'_1}^{\mathbf{d}'_1}$ that involves only arrows of x_j -height $\geq h$, as desired. $\square$

Corollary 8.4.2. *With the hypotheses of Lemma 8.4.1, suppose given two x_k -standard arrows $c_{j_1}^{\mathbf{d}_1}$ and $c_{j_2}^{\mathbf{d}_2}$ for β_t , and let $c_{j'_1}^{\mathbf{d}'_1}$, $c_{j'_2}^{\mathbf{d}'_2}$ be the associated liftings to β . If $c_{j'_1}^{\mathbf{d}'_1} \sim c_{j'_2}^{\mathbf{d}'_2}$, then $c_{j_1}^{\mathbf{d}_1} \sim c_{j_2}^{\mathbf{d}_2}$. Furthermore, if $c_{j'_1}^{\mathbf{d}'_1}$ can be advanced, then $c_{j_1}^{\mathbf{d}_1}$ can be advanced.*

Proof: Since $c_{j'_1}^{\mathbf{d}'_1}$ and $c_{j'_2}^{\mathbf{d}'_2}$ are x_k -standard and have x_j -height $\geq h$, Lemma 8.4.1 implies that there is a translation path from $c_{j'_1}^{\mathbf{d}'_1}$ to $c_{j'_2}^{\mathbf{d}'_2}$ consisting entirely of arrows of x_j -height $\geq h$. But then this path descends to give the translation equivalence of $c_{j_1}^{\mathbf{d}_1}$ and $c_{j_2}^{\mathbf{d}_2}$, which proves the first assertion.

Note that the second assertion is trivially true if

$$x_k\text{-degree}(\mathbf{x}^{\mathbf{d}_1}) > w_k^{(t)},$$

for then we can advance $c_{j_1}^{\mathbf{d}_1}$ by moving its tail in degree-decreasing steps toward the minimal generator $x_k^{w_k^{(t)}}$. It therefore remains to show that we can advance $c_{j_1}^{\mathbf{d}_1}$ provided that its lifting $c_{j'_1}^{\mathbf{d}'_1}$ can be advanced and

$$x_k\text{-degree}(\mathbf{x}^{\mathbf{d}'_1}) = x_k\text{-degree}(\mathbf{x}^{\mathbf{d}_1}) \leq w_k^{(t)}.$$

Since we can advance $c_{j'_1}^{\mathbf{d}'_1}$, we can translate it to an arrow $c_{j'_3}^{\mathbf{d}'_3}$ such that

$$\mathbf{x}^{\mathbf{d}'_3}/x_k \in I_\beta \text{ and } x_k\text{-degree}(\mathbf{x}^{\mathbf{d}'_3}) = x_k\text{-degree}(\mathbf{x}^{\mathbf{d}'_1}).$$

Then there is a minimal generator m of I_β such that

$$m \mid (\mathbf{x}^{\mathbf{d}'_3}/x_k) \text{ and } x_k\text{-degree}(m) > 0,$$

where the inequality follows from (15).

If $x_j\text{-degree}(m) < h$, then hypothesis (24) implies that

$$\begin{aligned} x_k\text{-degree}(m) \geq w_k^{(t)} &\Rightarrow x_k\text{-degree}(\mathbf{x}^{\mathbf{d}'_3}) > w_k^{(t)} \\ &\Rightarrow x_k\text{-degree}(\mathbf{x}^{\mathbf{d}'_1}) > w_k^{(t)}, \end{aligned}$$

which is a contradiction. We therefore have that

$$x_j\text{-degree}(m) \geq h \Rightarrow x_j\text{-degree}(\mathbf{x}^{\mathbf{d}'_3}) \geq h;$$

whence, Lemma 8.4.1 ensures that there is a translation path from $c_{j'_1}^{\mathbf{d}'_1}$ to $c_{j'_3}^{\mathbf{d}'_3}$ consisting entirely of arrows of x_j -height $\geq h$, but this path then descends to advance $c_{j_1}^{\mathbf{d}_1}$, which proves the second assertion. $\square$

Corollary 8.4.3. *Again with the hypotheses of Lemma 8.4.1, we have that the lifting, to β , of a standard x_k -sub-bunch $\mathcal{S}^{(t)}(k)$ of arrows for β_t , has maximal rank (mod M^2); furthermore, the lifted arrows cannot be advanced.*

Proof: It suffices, by Theorem 3.2.1, to show that for any two distinct arrows $c_{j_1}^{d_1}$, $c_{j_2}^{d_2} \in \mathcal{S}^{(t)}(k)$, with liftings $c_{j_1'}^{d_1'}$ and $c_{j_2'}^{d_2'}$, we have that $c_{j_1'}^{d_1'} \not\sim 0$ and $c_{j_1'}^{d_1'} \not\sim c_{j_2'}^{d_2'}$. Arguing by contradiction, suppose that $c_{j_1'}^{d_1'} \sim c_{j_2'}^{d_2'}$. Then Corollary 8.4.2 implies that $c_{j_1}^{d_1} \sim c_{j_2}^{d_2}$, which contradicts the hypothesis that $\mathcal{S}^{(t)}(k)$ is an x_k -standard sub-bunch. To prove that $c_{j_1'}^{d_1'} \not\sim 0$, it suffices to prove, more generally, that $c_{j_1'}^{d_1'}$ cannot be advanced. However, if this were false, then Corollary 8.4.2 would yield that $c_{j_1}^{d_1}$ can be advanced, which would again contradict the hypothesis that $\mathcal{S}^{(t)}(k)$ is an x_k -standard sub-bunch. $\square$

8.5. **x_k -sub-bunches of arrows for β and β_t , $x_k \neq x_j$.** We are now ready to prove

Lemma 8.5.1. *Let β_t be the x_j -truncation of β at height h , let $x_k \neq x_j$, and suppose that the specialization (24) of Hypothesis 8.1.1 to x_k holds. Then there exists a near-standard x_k -sub-bunch $\mathcal{S}'(k)$ of arrows for β that contains the lifting of a standard x_k -sub-bunch $\mathcal{S}^{(t)}(k)$ of arrows for β_t .*

Proof: We begin by constructing a standard x_k -sub-bunch $\mathcal{S}(k)$ for β as in the proof of Theorem 4.1.3. With $r(k, v)$ defined as in (23), we let

$$\mathcal{S}'_1(k) = \{c_{j_1}^{d_1} \in \mathcal{S}(k) \mid x_k\text{-deg}(\mathbf{x}^{j_1}) > r(k, v), \text{ where } v = x_k\text{-offset}(c_{j_1}^{d_1})\};$$

it is evident that $\mathcal{S}'_1$ is a maximal rank (mod M^2) set of x_k -standard arrows that cannot be advanced. It follows from Lemma 8.3.1 that the x_j -height of an arrow $c_{j_1}^{d_1} \in \mathcal{S}'_1(k)$ is $< h$. Furthermore, the cardinality of this set is

$$|\mathcal{S}'_1(k)| = |\beta| - |\beta_t| = n - n_t,$$

because shadow promotion does not change the x_k -heights of the heads of the promoted arrows (recall Equation (17)), and the arrows in the original sets $\mathcal{S}_0(k, v)$ with heads of height $\leq r(k, v)$ are precisely the arrows with heads in β_t .

We next construct a standard x_k -sub-bunch $\mathcal{S}^{(t)}(k)$ for β_t , and denote the lifting of this set to β by $\mathcal{S}'_2(k)$; this set consists of

$$|\mathcal{S}'_2(k)| = |\beta_t| = n_t$$

x_k -standard arrows $c_{j_2}^{d_2}$ having x_j -height $\geq h$. Corollary 8.4.3 implies that $\mathcal{S}'_2(k)$ has maximal rank (mod M^2), and that its arrows cannot be advanced. By Lemma 7.2.1, we know that the tail $\mathbf{x}^{d_2}$ is a minimal generator of I_β provided that its x_j -degree is $> h$; however, if the x_j -degree $= h$, we only know that $\mathbf{x}^{d_2}$ is an x_j -multiple of a minimal generator; therefore, some of the arrows in $\mathcal{S}'_2(k)$ may not be minimal.

The desired set is

$$\mathcal{S}'(k) = \mathcal{S}'_1(k) \cup \mathcal{S}'_2(k).$$

By comparing x_j -heights of arrows, we see that the two sets in the union do not overlap; therefore,

$$|\mathcal{S}'(k)| = |\mathcal{S}'_1(k)| + |\mathcal{S}'_2(k)| = (n - n_t) + n_t = n = |\beta|.$$

The arrows in $\mathcal{S}'(k)$ are x_k -standard arrows that cannot be advanced, but need not be minimal. To show that $\mathcal{S}'(k)$ is a near-standard x_k -sub-bunch, it remains to show that it has maximal rank (mod M^2). Since $\mathcal{S}'_1(k)$ and $\mathcal{S}'_2(k)$ each have maximal rank (mod M^2) and consist of x_k -unadvanceable arrows, it suffices (by Theorem 3.2.1) to prove that no arrow in $\mathcal{S}'_1(k)$ can be translated to an arrow in $\mathcal{S}'_2(k)$. So let $c_{j_1}^{d_1} \in \mathcal{S}'_1(k)$ have x_k -offset v . By definition we have that $x_k\text{-degree}(\mathbf{x}^{j_1}) > r(k, v)$. If this arrow could be translated to an arrow $c_{j_2}^{d_2}$ of x_j -height $\geq h$, then Lemma 8.3.1 would yield that $x_k\text{-degree}(\mathbf{x}^{j_2}) \leq r(k, v)$, implying that $c_{j_1}^{d_1}$ can be advanced, which is a contradiction. Therefore, $\mathcal{S}'(k)$ is a near-standard x_k -sub-bunch that contains the lifting of a standard x_k -sub-bunch for β_t , as desired. $\square$

8.6. Main theorem on truncations.

Theorem 8.6.1. *Let β be a smooth basis set, and β_t the x_j -truncation of β at height h . If in addition Hypothesis 8.1.1 holds, then β_t is a smooth basis set.*

Proof: To show that β_t is a smooth basis set, it suffices to prove that it satisfies conditions (a) and (b) of Theorem 5.1.1. Theorem 7.4.1 ensures that β_t satisfies condition (a) — that every non-standard arrow is translation-equivalent to 0 — because β , smooth by hypothesis, has this property; therefore, it remains to show that β_t satisfies condition (b).

Let $\mathcal{S}(j)$ be a standard x_j -sub-bunch of arrows for β , and $\mathcal{S}^{(t)}(j)$ the associated x_j -sub-bunch for β_t whose lifting to β lies in $\mathcal{S}(j)$, as in Lemma 8.2.1. For each variable $x_k \neq x_j$, let $\mathcal{S}'(k)$ be a near-standard x_k -sub-bunch of arrows for β constructed as in Lemma 8.5.1, and $\mathcal{S}^{(t)}(k)$ the associated x_k -sub-bunch for β_t whose lifting to β lies in $\mathcal{S}'(k)$. Taking unions, we obtain a near-standard bunch $\mathcal{S}'$ for β that contains the lifting of a standard bunch $\mathcal{S}^{(t)}$ for β_t . To complete the proof, it suffices to show that if $c_j^d \not\sim 0$ is a standard arrow for β_t , then $c_j^d \sim c_{j_1}^{d_1} \in \mathcal{S}^{(t)}$.

By assertion (c) of Lemma 4.2.1, we may assume that c_j^d is a minimal x_i -standard arrow for β_t that cannot be advanced. Consider the lifting $c_{j'}^{d'}$ of c_j^d to β . We have that $c_{j'}^{d'}$ cannot be advanced: Indeed, if $x_i = x_j$, then one checks easily that a translation path advancing $c_{j'}^{d'}$ would descend to a translation path advancing c_j^d , a contradiction. The same contradiction arises in case $x_i = x_k \neq x_j$ by Corollary 8.4.2. Since β is assumed smooth, we know that

$$c_{j'}^{d'} \sim c_{j_1}^{d_1} \in \mathcal{S}'.$$

Furthermore, since neither of the arrows $c_{j'}^{d'}$, $c_{j_1}^{d_1}$ can be advanced, we see that these arrows must have the same x_i -height. If $x_i = x_j$, we obtain that $c_{j_1}^{d_1}$ lies in the lifting

of $\mathcal{S}^{(t)}(j)$, and the translation path from $c_{j'}^{\mathbf{d}'}$ to $c_{j_1}^{\mathbf{d}'_1}$ descends to yield

$$c_j^{\mathbf{d}} \sim c_{j_1}^{\mathbf{d}_1} \in \mathcal{S}^{(t)}(j).$$

If $x_i = x_k \neq x_j$, we obtain a similar conclusion as follows: recalling that $\mathcal{S}'_2(k) \subseteq \mathcal{S}'(k)$ denotes the lifting of the x_k -standard sub-bunch $\mathcal{S}^{(t)}(k)$ (in the notation of Theorem 8.5.1), we *claim* that

$$(25) \quad c_{j'_1}^{\mathbf{d}'_1} \in \mathcal{S}'_2(k) \subseteq \mathcal{S}'(k).$$

Proof of claim: The lifted arrow $c_{j'}^{\mathbf{d}'}$ has x_j -height $\geq h$; therefore, by Lemma 8.3.1, we have that

$$x_k\text{-degree}(\mathbf{x}^{j'}) \leq r(k, v),$$

where $v = x_k\text{-offset}(c_{j'}^{\mathbf{d}'})$. Since the translation-equivalent arrow $c_{j'_1}^{\mathbf{d}'_1} \in \mathcal{S}'(k)$ has the same x_k -height and x_k -offset, it satisfies

$$x_k\text{-degree}(\mathbf{x}^{j'_1}) = x_k\text{-degree}(\mathbf{x}^{j'}) \leq r(k, v).$$

The claim follows immediately, because $\mathcal{S}'(k) = \mathcal{S}'_1(k) \cup \mathcal{S}'_2(k)$, and by definition the arrows $c_{j_*}^{\mathbf{d}^*} \in \mathcal{S}'_1(k)$ of x_k -offset v satisfy

$$x_k\text{-degree}(\mathbf{x}^{j_*}) > r(k, v).$$

The claim (25) implies that the arrow $c_{j'_1}^{\mathbf{d}'_1}$ is the lifting of an arrow $c_{j_1}^{\mathbf{d}_1} \in \mathcal{S}^{(t)}(k)$. Corollary 8.4.2 now yields that $c_j^{\mathbf{d}} \sim c_{j_1}^{\mathbf{d}_1}$, and the proof is complete. $\square$

9. ADDITION OF BOXES TO BASIS SETS

In this section we explore another way to construct a smooth basis set from a given smooth basis set, by “adding a box” in an appropriate way. The undoing of this operation (that is, the removal of the added box) is accomplished by a truncation.

9.1. Definition of box addition. Suppose that β is an arbitrary basis set of monomials in the variables $x_1, \dots, x_r$; recall that w_i denotes the x_i -width of β for $1 \leq i \leq r$. Choose one of the variables, say x_j , and an integer $h \geq 1$, and form the set

$$x_j^h \cdot \beta = \{x_j^h \cdot m \mid m \in \beta\},$$

which can be viewed geometrically as the translation of β a distance of h steps in the positive x_j -direction. Then, for each variable $x_k \neq x_j$, choose an integer $w'_k \geq w_k$, and form the box (20)

$$(26) \quad \mathcal{B} = \mathcal{B}(w'_1, w'_2, \dots, w'_{j-1}, h, w'_{j+1}, \dots, w'_r).$$

We then set

$$(27) \quad \beta' = (x_j^h \cdot \beta) \cup \mathcal{B},$$




FIGURE 5. The basis set on the right is obtained by adding a box in the x_2 -direction to the basis set on the left.

and say that β' is obtained by **adding the box $\mathcal{B}$ in the x_j -direction to β** (see Figure 5).

Lemma 9.1.1. *Let β' be the set of monomials obtained by adding the box $\mathcal{B}$ (26) to the basis set β in the x_j -direction, as in (27). Then*

- (a) β' is a basis set.
- (b) β is the x_j -truncation of β' at height h .

Proof: To prove that β' is a basis set, we must show that if m_1 and m_2 are monomials such that $m_1 \in \beta'$ and $m_2 \mid m_1$, then $m_2 \in \beta'$. If $m_1 \in \mathcal{B}$, then it is clear that $m_2 \in \mathcal{B} \subseteq \beta'$. If $m_1 \notin \mathcal{B}$, then $m_1/x_j^h \in \beta$. Since $m_2 \mid m_1$, it is clear that

$$x_k \neq x_j \Rightarrow x_k\text{-degree}(m_2) \leq x_k\text{-degree}(m_1/x_j^h) < w_k \leq w'_k;$$

therefore,

$$x_j\text{-degree}(m_2) < h \Rightarrow m_2 \in \mathcal{B} \subseteq \beta',$$

and

$$\begin{aligned} x_j\text{-deg}(m_2) \geq h &\Rightarrow (m_2/x_j^h) \mid (m_1/x_j^h) \\ &\Rightarrow m_2/x_j^h \in \beta \\ &\Rightarrow m_2 \in x_j^h \cdot \beta \subseteq \beta'. \end{aligned}$$

This completes the proof that β' is a basis set, and the second statement follows easily from the definitions. $\square$

9.2. Minimal generators of $I_{\beta'}$. Let $\beta, \beta', \mathcal{B}$, etc., be as above. Since $\beta = \beta'_t$ is the x_j -truncation of β' at height h , we have that

$$w'_j = w_j + h$$

is the x_j -width of β' , by assertion (d) of Lemma 7.2.1. We then have the following

Lemma 9.2.1. *For each i , $1 \leq i \leq r$, the x_i -width of β' is w'_i ; that is, $x_i^{w'_i}$ is a minimal generator of $I_{\beta'}$. Furthermore, for all minimal generators m of $I_{\beta'}$ we have that*

$$x_j\text{-degree}(m) < h \Rightarrow m = x_k^{w'_k} \text{ for some } x_k \neq x_j;$$

in particular, Hypothesis 8.1.1 holds for β' and its truncation β .

Proof: For all variables $x_i = x_k \neq x_j$, we have that

$$x_k^{w'_k} \in I_{\beta'} \subseteq I_{\mathcal{B}},$$

and $x_k^{w'_k}$ is a minimal generator of $I_{\mathcal{B}}$, by Lemma 6.5.1; therefore, $x_k^{w'_k}$ is a minimal generator of $I_{\beta'}$. For $x_i = x_j$, we have already observed that w'_j is the x_j -width of β' , so the first assertion holds. Suppose now that m is a minimal generator of $I_{\beta'}$ such that $x_j\text{-degree}(m) < h$. A moment's reflection shows that in fact m must be a minimal generator of the monomial ideal $I_{\mathcal{B}}$; the second assertion then follows at once from Lemma 6.5.1. $\square$

In addition, recall that Lemma 7.2.1 further describes the relationship between the minimal generators of $I_{\beta'}$ and the minimal generators of I_{β} , since β is the x_j -truncation of β' at height h .

9.3. Main theorem on box additions. Box addition is a convenient tool for building smooth basis sets, as the following result suggests:

Theorem 9.3.1. *Let β be a basis set, and let β' be obtained by adding the box $\mathcal{B}$ (26) in the x_j -direction. Then:*

$$\beta \text{ is smooth} \Leftrightarrow \beta' \text{ is smooth}.$$

Proof: ($\Leftarrow$): Since β' is smooth and Hypothesis 8.1.1 holds, by Lemma 9.2.1, Theorem 8.6.1 implies that the truncation $\beta'_t = \beta$ is smooth.

($\Rightarrow$): Given that β is a smooth basis set, we must prove that β' is a smooth basis set. To do this we will use Theorem 5.1.1: we must show (for β') that every non-standard arrow is translation-equivalent to 0, and that there exists a near-standard bunch of arrows $\mathcal{S}'$ such that if $c_{\mathbf{j}}^{\mathbf{d}} \not\sim 0$ is an x_i -standard arrow, then there is an x_i -standard arrow $c_{\mathbf{j}'}^{\mathbf{d}'} \in \mathcal{S}'$ such that $c_{\mathbf{j}}^{\mathbf{d}} \sim c_{\mathbf{j}'}^{\mathbf{d}'}$.

First, let $c_{\mathbf{j}}^{\mathbf{d}}$ be a non-standard arrow, which we can assume is minimal by Lemma 4.1.2. Since the arrow is non-standard we know that the vector $\mathbf{j} - \mathbf{d}$ has negative components in at least two variable directions, say x_{i_1} and x_{i_2} . Note that the tail of the arrow must be a minimal generator of $I_{\beta'}$ that has x_j -degree $\geq h$, since the minimal generators of β' with x_j -degree $< h$ are the corner monomials $x_k^{w'_k}$ (Lemma 9.2.1), and a non-standard arrow cannot have a corner monomial as its tail.

Suppose first that the head of the arrow $\mathbf{x}^{\mathbf{j}} \in \mathcal{B}$. By translating the arrow in the increasing x_{i_2} -direction, one eventually reaches an arrow $c_{\mathbf{j}_1}^{\mathbf{d}_1} \sim c_{\mathbf{j}}^{\mathbf{d}}$ such that

$$x_{i_2}\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) = w'_{i_2} - 1 \Rightarrow x_{i_2}\text{-degree}(\mathbf{x}^{\mathbf{d}_1}) \geq w'_{i_2};$$

therefore, $c_{\mathbf{j}_1}^{\mathbf{d}_1}$ can be translated in the direction of decreasing x_{i_1} -degree until the head exits the first orthant; whence, $c_{\mathbf{j}}^{\mathbf{d}} \sim 0$.

If $\mathbf{x}^j \notin \mathcal{B}$, then $\mathbf{x}^j \in x_j^h \cdot \beta$, and the arrow $c_j^{\mathbf{d}}$ descends to $c_{j_0}^{\mathbf{d}_0}$. Since β is assumed smooth, we have that $c_{j_0}^{\mathbf{d}_0} \sim 0$. If the associated translation path causes the head to cross the hyperplane (x_k -degree = 0) for any $x_k \neq x_j$, then we can lift the translation path to β' to obtain that $c_j^{\mathbf{d}} \sim 0$. If the head crosses the hyperplane (x_j -degree = 0), then the lifted path shows that $c_j^{\mathbf{d}} \sim c_{j_2}^{\mathbf{d}_2}$ with $\mathbf{x}^{j_2} \in \mathcal{B}$, and $c_{j_2}^{\mathbf{d}_2} \sim 0$ as before.

Now let $\mathcal{S}'$ be the near-standard bunch of arrows for β' that was constructed in the proof of Theorem 8.6.1 (based on Lemmas 8.2.1 and 8.5.1); recall that $\mathcal{S}'$ contains the lifting of a standard sub-bunch $\mathcal{S}^{(t)}$ for $\beta = \beta'_t$. Suppose that $c_j^{\mathbf{d}} \not\sim 0$ is an x_i -standard arrow; we can assume that this arrow is minimal and cannot be advanced by assertion (c) of Lemma 4.2.1. We must show that $c_j^{\mathbf{d}}$ is translation-equivalent to an arrow in $\mathcal{S}'(i)$.

Suppose first that $x_i = x_j$, so that $\mathcal{S}'(i) = \mathcal{S}(j)$ is a standard x_j -sub-bunch. If $\mathbf{x}^j \notin \mathcal{B}$, then the arrow $c_j^{\mathbf{d}}$ is the lifting of a minimal standard arrow $c_{j_0}^{\mathbf{d}_0}$ for β . Since β is smooth, we can find a minimal standard arrow $c_{j'_0}^{\mathbf{d}'_0} \in \mathcal{S}^{(t)}(j)$ such that $c_{j_0}^{\mathbf{d}_0} \sim c_{j'_0}^{\mathbf{d}'_0}$. Lifting the translation path, we find that $c_j^{\mathbf{d}} \sim c_{j'}^{\mathbf{d}'} \in \mathcal{S}'(j)$.

If $\mathbf{x}^j \in \mathcal{B}$, then let $c_{j'}^{\mathbf{d}'} \in \mathcal{S}'(j)$ be the unique arrow such that

$$x_j\text{-degree}(\mathbf{x}^{j'}) = x_j\text{-degree}(\mathbf{x}^j) \quad \text{and} \quad x_j\text{-offset}(c_{j'}^{\mathbf{d}'}) = x_j\text{-offset}(c_j^{\mathbf{d}}),$$

the existence of which is ensured by Corollary 4.6.1. Suppose that the lengths of these two arrows differ; in other words, suppose that

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{d}}) \neq x_j\text{-degree}(\mathbf{x}^{\mathbf{d}'}).$$

Recalling Lemma 6.5.2, we let

$$\mathbf{x}^{j_1} = \text{lcm}(\mathbf{x}^j, \mathbf{x}^{j'}) \in \mathcal{B} \subseteq \beta'.$$

It is clear that we can translate both $c_j^{\mathbf{d}}$ and $c_{j'}^{\mathbf{d}'}$ by degree-increasing steps (excluding the x_j -direction) to arrows $c_{j_1}^{\mathbf{d}_1}$ and $c_{j_1}^{\mathbf{d}'_1}$, respectively. The tails $\mathbf{x}^{\mathbf{d}_1}$ and $\mathbf{x}^{\mathbf{d}'_1}$ differ only in x_j -degree; it follows that the arrow corresponding to the tail of larger x_j -degree can be advanced, which is a contradiction, since neither $c_j^{\mathbf{d}}$ nor $c_{j'}^{\mathbf{d}'}$ can be advanced. We therefore have that

$$c_{j_1}^{\mathbf{d}_1} = c_{j_1}^{\mathbf{d}'_1} \Rightarrow c_j^{\mathbf{d}} \sim c_{j'}^{\mathbf{d}'} \in \mathcal{S}'(j),$$

as desired.

Finally, we have to consider the case in which $c_j^{\mathbf{d}}$ is a minimal x_k -standard arrow that cannot be advanced, where $x_k \neq x_j$. Let v denote the x_k -offset of $c_j^{\mathbf{d}}$. From Lemma 9.2.1, we know that the tail $\mathbf{x}^{\mathbf{d}}$ is either equal to the corner monomial $x_k^{w'_k}$ or else $x_j\text{-degree}(\mathbf{x}^{\mathbf{d}}) \geq h$.

In case $\mathbf{x}^{\mathbf{d}} = x_k^{w'_k}$, if the head

$$\mathbf{x}^j \notin \beta = \beta'_t,$$




FIGURE 6. A sequence of compound boxes, beginning on the left with a box.

then

$$x_k\text{-degree}(\mathbf{x}^j) > r(k, v) \Rightarrow c_j^{\mathbf{d}} \in \mathcal{S}'_1(k) \subseteq \mathcal{S}'(k),$$

in the notation of Lemma 8.5.1. If the head

$$\mathbf{x}^j \in \beta = \beta'_t,$$

then, since $c_j^{\mathbf{d}}$ cannot be advanced, but *can* be translated h steps in the direction of increasing x_j -degree to $c_{j'}^{\mathbf{d}'}$, we must have that $w'_k = w_k$. This in turn implies that $c_j^{\mathbf{d}}$, considered as an arrow for the truncation β , lies in the sub-bunch $\mathcal{S}^{(t)}(k)$; therefore,

$$c_j^{\mathbf{d}} \sim c_{j'}^{\mathbf{d}'} \in \mathcal{S}'_2(k) \subseteq \mathcal{S}'(k).$$

In case $x_j\text{-degree}(\mathbf{x}^{\mathbf{d}}) \geq h$, we can descend the arrow to $c_{j_0}^{\mathbf{d}_0}$; then, because β is assumed smooth, we know that $c_{j_0}^{\mathbf{d}_0} \sim c_{j'_0}^{\mathbf{d}'_0} \in \mathcal{S}^{(t)}(k)$, and the translation path lifts to yield $c_j^{\mathbf{d}} \sim c_{j'}^{\mathbf{d}'} \in \mathcal{S}'(k)$. This completes the proof of the theorem. $\square$

9.4. Compound boxes. By a **compound box**, we mean a basis set β that is constructed by starting with a box, and then performing a finite sequence of box additions in various variable directions; Figure 6 illustrates the idea. Note that a box $\mathcal{B}$ is a compound box, since it can be generated by starting with itself and performing a sequence of box additions of length 0.

Since the starting point (a box) is smooth, by Proposition 6.5.3, and adding a box to a smooth basis set yields a smooth basis set, by Theorem 9.3.1, we obtain the following corollary by induction:

Corollary 9.4.1. *If β is a compound box, then β is a smooth basis set.* $\square$

9.5. Example: Basis sets in two variables. It is easy to verify (see Figure 7) that *every* basis set β in two variables is a compound box; whence, Corollary 9.4.1 yields the following

Corollary 9.5.1. *Every basis set β in two variables is smooth.* $\square$

Haiman's lovely proof of this result (part of the proof of [6, Proposition 2.4]) introduced the idea of arrow translation, and inspired the present paper. Note that the $\mathbf{k}$ -basis of M/M^2 that he obtains is slightly different from ours; his basis arrows




FIGURE 7. Every basis set in two variables is a compound box.

are typically not *minimal* standard arrows. From the smoothness of the points t_β , Haiman deduces that $\mathbf{H}^n$ is everywhere nonsingular and irreducible (facts first proved by Fogarty [1]). Corollary 9.4.1 can be viewed as a generalization of the two-variable smoothness phenomenon to higher dimensions.

9.6. Example: The lexicographic point. The “lexicographic point” of $\text{Hilb}_{\mathbb{P}^r}^{p(z)}$ is the point corresponding to the unique saturated lexicographic ideal L such that $\mathbf{k}[X_0, \dots, X_r]/L$ has Hilbert polynomial $p(z)$. A. Reeves and M. Stillman prove in general that the lexicographic point is a smooth point [12]. In the case of a constant Hilbert polynomial $p(z) = n$, one checks that

$$L = (X_0, X_1, \dots, X_{r-2}, X_{r-1}^n).$$

Dehomogenizing with respect to the variable X_r , we obtain the ideal $I_\beta \subseteq \mathbf{k}[x_0, \dots, x_{r-1}]$, where

$$\beta = \{1, x_{r-1}, x_{r-1}^2, \dots, x_{r-1}^{n-1}\}.$$

This is clearly a smooth basis set, since it is a special case of a box.

9.7. Example: $\beta = \{1, x_1, x_2, x_1x_2, x_3\}$. In Example 5.3 we considered the basis set

$$\{1, x_1, x_2, x_3\},$$

which is non-smooth. If we add the monomial x_1x_2 , we obtain a compound box β (see Figure 8); we will verify “by hand” that β is smooth.

The minimal generators of I_β are the monomials

$$x_1^2, x_2^2, x_1x_3, x_2x_3, x_3^2;$$

whence, one has $5 \cdot 5 = 25$ minimal arrows, and these span M/M^2 by Lemma 4.1.2. However, inspecting the minimal arrows, we find that:

- Four are non-standard arrows, all of which are translation-equivalent to 0: $c_{(0,0,0)}^{(1,0,1)}$, $c_{(0,1,0)}^{(1,0,1)}$, $c_{(0,0,0)}^{(0,1,1)}$, and $c_{(1,0,0)}^{(0,1,1)}$.
- Five are standard arrows that are translation-equivalent to 0: $c_{(0,0,0)}^{(2,0,0)}$, $c_{(0,0,0)}^{(0,2,0)}$, $c_{(0,0,0)}^{(0,0,2)}$, $c_{(1,0,0)}^{(0,0,2)}$, and $c_{(0,1,0)}^{(0,0,2)}$.
- Two are standard arrows that are not translation-equivalent to 0, but are translation-equivalent to each other: $c_{(1,0,0)}^{(1,0,1)} \sim c_{(0,1,0)}^{(0,1,1)}$.




FIGURE 8. The compound box $\beta = \{1, x_1, x_2, x_1x_2, x_3\}$, with the minimal generators of I_β shown in boldface. Note, for example, that the arrows $c_{(0,1,0)}^{(1,0,1)}$ and $c_{(1,0,0)}^{(0,0,2)}$ are translation-equivalent to 0, and $c_{(1,0,0)}^{(1,0,1)} \sim c_{(0,1,0)}^{(0,1,1)}$.

This means that there are at most 15 non-trivial translation-equivalence classes of arrows available to span M/M^2 , but $r \cdot n = 3 \cdot 5 = 15$ is a lower bound on the k -dimension of the cotangent space, by Proposition 2.4.1. It follows that M/M^2 has k -dimension 15; whence, β is smooth.

10. SMOOTH BASIS SETS IN THREE VARIABLES ARE COMPOUND BOXES

The main goal of this section is to prove Theorem 10.3.1, which states that a basis set β in three variables is smooth if and only if β is a compound box. In the next section we will show by example that smooth basis sets in four or more variables need not be compound boxes.

10.1. The main lemma. Let β be a basis set in the variables x_1, x_2, x_3 , and I_β the associated monomial ideal. For

$$(28) \quad (i, j) \in \{(1, 2), (1, 3), (2, 3)\},$$

we write $G(i, j)$ for the set of minimal generators of I_β that involve only variables in the set $\{x_i, x_j\}$. For example, we always have that

$$x_i^{w_i} \in G(i, j) \quad \text{and} \quad x_j^{w_j} \in G(i, j),$$

so the number of elements

$$|G(i, j)| \geq 2.$$

Lemma 10.1.1. *If $|G(i, j)| > 2$ for all of the ordered pairs (i, j) in (28), then there exists a non-standard arrow for β that is not translation-equivalent to 0; consequently, β is not smooth by Theorem 5.1.1.*

10.2. Proof of Lemma 10.1.1. By the hypothesis, we may choose

$$(29) \quad \begin{aligned} m_{(1,3)} &= x_1^a x_3^b \in G(1, 3), \\ m_{(2,3)} &= x_2^c x_3^d \in G(2, 3), \\ m_{(1,2)} &= x_1^e x_2^f \in G(1, 2) \end{aligned}$$




FIGURE 9. The two cases considered in the proof of Lemma 10.1.1. In the left-hand figure, the monomial $m_{(1,3)}$ is dominant; the associated rigid non-standard arrow is also illustrated in Figure 1. In the right-hand figure, none of the monomials $m_{(i,j)}$ are dominant. The associated non-standard arrow is not translation-equivalent to 0.

such that all the exponents are positive and $m_{(1,3)}$ (resp. $m_{(2,3)}$) has maximal x_1 -degree (resp. maximal x_2 -degree) subject to the stated constraints; note that this implies that $m_{(1,3)}$ (resp. $m_{(2,3)}$) has minimal x_3 -degree subject to the stated constraints. We proceed to construct the desired non-standard arrow; there are two cases (see Figure 9):

10.2.1. *Case 1: one of the monomials in (29) is dominant.* We say that $m_{(i,j)}$ is **dominant** provided that the degree of each of its constituent variables is greater than or equal to the degree of the same variable in the other monomial in which it appears. For example, $m_{(1,3)}$ is dominant provided that (as shown in the left-hand portion of Figure 9)

$$a \geq e \text{ and } b \geq d.$$

In this case, let $g = \max(c, f)$, and note that x_2^g is a basis monomial. Starting at x_2^g , we can move to a maximal basis monomial $m^* = x_1^p x_2^q x_3^r$ by a sequence of degree-increasing steps. Since $x_1^e x_2^f$ is a minimal generator of the ideal, we know that $p < e$. Similarly, since $x_2^c x_3^d$ is a minimal generator, we have that $r < d$. Then the rigid arrow A with tail $m_{(1,3)}$ and head m^* is also non-standard, since its vector

$$(p, q, r) - (a, 0, b) = (p - a, q, r - b)$$

has the first and third coordinates negative:

$$\begin{aligned} p - a &< e - a \leq a - a = 0, \\ r - b &< d - b \leq b - b = 0. \end{aligned}$$

Since A is rigid, it is not translation-equivalent to 0, as desired.

10.2.2. *Case 2: None of the monomials in (29) is dominant.* Writing out what this condition says, we find that:

$$\begin{aligned}
& (\neg(a \geq e \wedge b \geq d)) \wedge (\neg(c \geq f \wedge d \geq b)) \wedge (\neg(e \geq a \wedge f \geq c)) \\
& \quad \equiv \\
& (a < e \vee b < d) \wedge (c < f \vee d < b) \wedge (e < a \vee f < c) \\
& \quad \equiv \\
& (a < e \wedge f < c \wedge d < b) \vee (b < d \wedge c < f \wedge e < a).
\end{aligned}$$

Suppose the first alternative in the last line holds; that is, suppose (as shown in the right-hand portion of Figure 9) that

$$a < e, \quad f < c, \quad \text{and} \quad d < b.$$

Note first of all that the monomial

$$m = x_2^{(w_2-1)} x_3^{(d-1)} \in \beta.$$

If not, there would exist a minimal generator m' of I_β such that

$$m' \mid m \Rightarrow m' \in G(2, 3).$$

It is clear that m' cannot equal either of the corner monomials $x_2^{w_2}, x_3^{w_3}$, so m' must involve both x_2 and x_3 nontrivially. However, the x_3 -degree of m' is $\leq d-1$, which contradicts our choice of $m_{(2,3)}$ as having minimal x_3 -degree d among the members of $G(2, 3)$ that involve two variables nontrivially.

Let $s > 0$ denote the minimal exponent such that

$$x_1^s \cdot m \notin \beta;$$

since

$$(x_1^e x_2^f = m_{(1,2)}) \mid (x_1^e \cdot m) \Rightarrow (x_1^e \cdot m) \notin \beta,$$

we have that

$$(30) \quad s \leq e < w_1,$$

where the second inequality holds because the minimal generator $x_1^{w_1}$ cannot divide the minimal generator $m_{(1,2)}$. Let

$$(31) \quad \alpha = \begin{cases} s-1, & \text{if } s \leq a, \\ a-1, & \text{if } s > a, \end{cases}$$

and form the arrow

$$A = c_{(\alpha, w_2-1, d-1)}^{(a, 0, b)}$$

with tail $m_{(1,3)}$ and head

$$m_h = x_1^\alpha x_2^{w_2-1} x_3^{d-1} \in \beta \quad (\text{since } \alpha < s).$$

Notice that A is non-standard, since its vector

$$(\alpha, w_2-1, d-1) - (a, 0, b) = (\alpha-a, w_2-1, (d-1)-b)$$

has negative first and third components (and non-negative second component w_2-1).

We assert that this arrow is not translation-equivalent to 0. If it were, the head would have to exit the first octant across either the hyperplane (x_1 -degree = 0) or the hyperplane (x_3 -degree = 0). In the former case, we would have to translate A to an arrow A' of the same x_1 -height ($= a$), but with tail divisible by a minimal generator $\tilde{m}$ of x_1 -degree $< a$. Since $\tilde{m}$ cannot divide $m_{(1,2)}$ and $m_{(1,3)}$, we must have that

$$(32) \quad \text{either } x_2\text{-degree}(\tilde{m}) > f \text{ or } x_3\text{-degree}(\tilde{m}) > b.$$

However, we have that A is “rigid” with respect to motion in the x_2 -direction; that is, neither A nor any of its translates A' can be moved in either the increasing or decreasing x_2 -direction. Indeed, an easy induction on the length of the path from A to A' shows that

$$x_2\text{-degree}(\text{head}(A')) = w_2 - 1 \text{ and } x_2\text{-degree}(\text{tail}(A')) = 0,$$

and clearly such an arrow cannot be translated in either the increasing or decreasing x_2 -degree directions. Furthermore, neither A nor any of its translates A' has x_3 -height greater than the initial value b , for we have already seen that the x_2 -degree of the head of A' is invariantly $w_2 - 1$, and if the x_3 -height of A' were to exceed b , then the x_3 -degree of the head would be $\geq d$, implying that the head would be divisible by $m_{(2,3)}$. Therefore, we see that it is impossible to translate A to A' with tail divisible by $\tilde{m}$ as in (32), so we cannot move the head of A across the hyperplane (x_1 -degree = 0).

The only remaining possibility is to translate A so that the head exits the first octant across the hyperplane (x_3 -degree = 0). This requires us to translate A to an arrow A' of the same x_3 -height ($= b$), but with tail divisible by a minimal generator $\hat{m}$ of x_3 -degree $< b$. Since the x_2 -height of A' is invariantly 0, we see that

$$x_2\text{-degree}(\hat{m}) = 0 \Rightarrow \hat{m} \in G(1, 3) \Rightarrow \hat{m} = x_1^{w_1},$$

where the last implication follows from our choice of $m_{(1,3)}$ as the monomial of minimal x_3 -degree in $G(1, 3)$ among those involving both x_1 and x_3 nontrivially. In view of the constraints on the motion of A , we see that we would have to be able to translate A a distance of $w_1 - a$ units in the positive x_1 -direction, but this motion would move the head to

$$x_1^{\alpha+(w_1-a)} x_2^{w_2-1} x_3^{d-1},$$

which lies outside of β , because (recalling (30) and (31))

$$\begin{aligned} s \leq a &\Rightarrow \alpha + (w_1 - a) = (s - 1) + (w_1 - a) \geq s, \text{ and} \\ s > a &\Rightarrow \alpha + (w_1 - a) = (a - 1) + (w_1 - a) = w_1 - 1 \geq s. \end{aligned}$$

The required translation is therefore impossible; whence, A is not translation-equivalent to 0, and the proof of Lemma 10.1.1 is complete. $\square$

10.3. The main theorem.

Theorem 10.3.1. *Let β be a basis set in the variables x_1, x_2, x_3 . Then*

$$\beta \text{ is smooth} \Leftrightarrow \beta \text{ is a compound box.}$$

Proof: ($\Leftarrow$): Immediate from Corollary 9.4.1.

($\Rightarrow$): We proceed by induction on $n = |\beta|$.

Base case: $n = 1$. The only basis set of cardinality 1 is the box

$$\{1\} = \mathcal{B}(1, 1, 1),$$

which is smooth by Proposition 6.5.3, and trivially a compound box.

Inductive step: Suppose that $|\beta| = n$, and that any smooth basis set of cardinality $< n$ is a compound box. Since β is by hypothesis smooth, Lemma 10.1.1 implies that for at least one of the pairs (i, j) in (28), we have that

$$G(i, j) = \{x_i^{w_i}, x_j^{w_j}\};$$

that is, no minimal generator of I_β exists that involves both x_i and x_j nontrivially, but does not involve the third variable x_k . Let h be the minimal x_k -degree among the minimal generators of I_β that *do* involve x_k . If $h = w_k$, then

$$\beta = \mathcal{B}(w_1, w_2, w_3)$$

is a (compound) box, as desired. If $h < w_k$, we let β_t be the x_k -truncation of β at height h (21). Then one sees easily that β is obtained from β_t by adding a box in the x_k -direction, as described in Section 9: more precisely, the added box has dimensions w_i in the x_i -direction, w_j in the x_j -direction, and h in the w_k -direction. It follows from Theorem 9.3.1 that β_t is smooth, since β is smooth by hypothesis. The induction hypothesis now implies that β_t is a compound box $\Rightarrow \beta$ is a compound box, and we are done. $\square$

More can be gleaned from the preceding proof: suppose that β is a basis set in the variables x_1, x_2, x_3 for which condition (a) of Theorem 5.1.1 holds; that is, every non-standard arrow for β is translation-equivalent to 0. Then Lemma 10.1.1 implies that for at least one of the pairs (i, j) in (28), we have that

$$G(i, j) = \{x_i^{w_i}, x_j^{w_j}\};$$

arguing as in the preceding proof, we then see that β is either a box or is the result of adding a box to a truncation β_t . In the former case, β is smooth, and in the latter case, β_t also satisfies condition (a) of Theorem 5.1.1, by Theorem 7.4.1; therefore, induction yields that β_t is smooth $\Rightarrow \beta$ is smooth, by Theorem 9.3.1. Whence:

Corollary 10.3.2. *Let β be a basis set in the variables x_1, x_2, x_3 . Then*

$$\beta \text{ is smooth} \Leftrightarrow \begin{cases} \text{condition (a) of Theorem 5.1.1 holds; that is, every} \\ \text{non-standard arrow for } \beta \text{ is translation-equivalent to 0.} \end{cases} \quad \square$$

As of this writing, I do not know if this result extends to higher dimensions.

11. THE UNION OF TWO BOXES

To end this paper, we study one more family of smooth basis sets that does not consist entirely of compound boxes (in four or more variables): basis sets that are unions of two boxes.

11.1. Notation. We will use the following notation throughout this section. Let

$$\begin{aligned}\mathcal{B}_1 &= \mathcal{B}(w_{1,1}, w_{2,1}, \dots, w_{r,1}), \\ \mathcal{B}_2 &= \mathcal{B}(w_{1,2}, w_{2,2}, \dots, w_{r,2})\end{aligned}$$

be two boxes (20) in the variables $x_1, \dots, x_r$, and let

$$(33) \quad \beta = \mathcal{B}_1 \cup \mathcal{B}_2.$$

One checks easily that β is a basis set. Lacking inspiration, we call β a **two-box union**.

11.2. Minimal generators of I_β . As usual, we write w_i for the x_i -width of β ; that is, $x_i^{w_i}$ is the corner (minimal) monomial divisible by x_i . The following result is clear:

Lemma 11.2.1. *The x_i -width w_i of the two-box union β (33) is given by*

$$w_i = \max(w_{i,1}, w_{i,2}), \quad 1 \leq i \leq r. \quad \square$$

We now write the set of variables as a union

$$\{x_1, x_2, \dots, x_r\} = V_1 \cup V_2 \cup V_3,$$

where

$$(34) \quad \begin{aligned}V_1 &= \{x_j \mid w_{j,1} > w_{j,2}\}, \\ V_2 &= \{x_k \mid w_{k,1} < w_{k,2}\}, \\ V_3 &= \{x_\ell \mid w_{\ell,1} = w_{\ell,2}\};\end{aligned}$$

henceforth, we will use the subscripts j , k , and ℓ to denote membership in V_1 , V_2 , and V_3 , respectively. One sees easily that

$$\begin{aligned}V_1 = \emptyset &\Rightarrow \mathcal{B}_1 \subseteq \mathcal{B}_2 \Rightarrow \beta = \mathcal{B}_2, \\ V_2 = \emptyset &\Rightarrow \mathcal{B}_2 \subseteq \mathcal{B}_1 \Rightarrow \beta = \mathcal{B}_1,\end{aligned}$$

so the most interesting case is when both V_1 and V_2 are non-empty.

We have the following

Lemma 11.2.2. *Let β be a two-box union (33). Then a minimal generator m of the monomial ideal I_β is either a corner monomial $x_i^{w_i}$ or a two-variable monomial of the form $x_j^{w_{j,2}} x_k^{w_{k,1}}$, with $x_j \in V_1$, $x_k \in V_2$ (34).*

Proof: Let

$$m = x_1^{s_1} x_2^{s_2} \dots x_r^{s_r}$$

be a minimal generator of I_β , and suppose that three or more of the exponents (say s_1 , s_2 , and s_3) are positive. Then each of the monomials

$$x_1^{s_1-1}x_2^{s_2}x_3^{s_3}\dots x_r^{s_r}, \quad x_1^{s_1}x_2^{s_2-1}x_3^{s_3}\dots x_r^{s_r}, \quad x_1^{s_1}x_2^{s_2}x_3^{s_3-1}\dots x_r^{s_r}$$

belong to β , which implies that two of these monomials must belong to the same box ($\mathcal{B}_1$ or $\mathcal{B}_2$). But then the least common multiple of the two monomials must also belong to this box (Lemma 6.5.2); that is, $m \in \beta$, which is a contradiction. We conclude that ≤ 2 of the exponents s_i can be positive. If only one of the exponents s_i is positive, then m is the corner monomial $x_i^{(s_i=w_i)}$. If two of the exponents (say s_1 and s_2) are positive, so that

$$m = x_1^{s_1}x_2^{s_2},$$

we again have that the monomials

$$x_1^{s_1-1}x_2^{s_2}, \quad x_1^{s_1}x_2^{s_2-1}$$

belong to β ; if both belonged to the same box, then we would arrive once more at the contradiction $m \in \beta$. Therefore, the latter two monomials belong to different boxes, say

$$x_1^{s_1-1}x_2^{s_2} \in \mathcal{B}_2 \setminus \mathcal{B}_1, \quad x_1^{s_1}x_2^{s_2-1} \in \mathcal{B}_1 \setminus \mathcal{B}_2;$$

consequently,

$$\begin{aligned} x_1^{s_1-1}x_2^{s_2} \in \mathcal{B}_2 &\Leftrightarrow s_1 - 1 < w_{1,2} \text{ and } s_2 < w_{2,2}, \\ x_1^{s_1}x_2^{s_2-1} \in \mathcal{B}_1 &\Leftrightarrow s_1 < w_{1,1} \text{ and } s_2 - 1 < w_{2,1}, \end{aligned}$$

and

$$\begin{aligned} x_1^{s_1-1}x_2^{s_2} \notin \mathcal{B}_1 &\Leftrightarrow s_1 - 1 \geq w_{1,1} \text{ or } s_2 \geq w_{2,1}, \\ x_1^{s_1}x_2^{s_2-1} \notin \mathcal{B}_2 &\Leftrightarrow s_1 \geq w_{1,2} \text{ or } s_2 - 1 \geq w_{2,2}. \end{aligned}$$

Note that

$$s_2 < w_{2,2} \Rightarrow s_2 - 1 < w_{2,2} \Rightarrow s_1 \geq w_{1,2};$$

whence,

$$(s_1 \geq w_{1,2} \text{ and } s_1 - 1 < w_{1,2}) \Rightarrow s_1 = w_{1,2},$$

and

$$w_{1,2} = s_1 < w_{1,1} \Rightarrow x_1 \in V_1,$$

as desired. A similar argument yields

$$s_2 = w_{2,1} \text{ and } x_2 \in V_2,$$

and the proof is complete. $\square$

11.3. Two-box unions are smooth basis sets. Retaining the notation of Subsections 11.1 and 11.2, we begin with the following

Lemma 11.3.1. *Let β be a two-box union, and let*

$$\mathbf{x}^{\mathbf{d}} = x_j^{w_{j,2}} x_k^{w_{k,1}}, \quad x_j \in V_1, \quad x_k \in V_2,$$

be a two-variable minimal generator of I_β , as in Lemma 11.2.2. If $c_j^{\mathbf{d}}$ is an x_j - (resp. x_k -) standard arrow for β , then the head of the arrow $\mathbf{x}^{\mathbf{j}} \in \mathcal{B}_2$ (resp. $\mathcal{B}_1$).

Proof: If $c_j^{\mathbf{d}}$ is an x_j -standard arrow, then we have that

$$x_k\text{-degree}(\mathbf{x}^{\mathbf{j}}) \geq x_k\text{-degree}(\mathbf{x}^{\mathbf{d}}) = w_{k,1} \Rightarrow \mathbf{x}^{\mathbf{j}} \notin \mathcal{B}_1.$$

Since $\mathbf{x}^{\mathbf{j}} \in \beta$, we must have that $\mathbf{x}^{\mathbf{j}} \in \mathcal{B}_2$, as asserted. A similar argument applies in the case that $c_j^{\mathbf{d}}$ is x_k -standard. $\square$

Theorem 11.3.2. *A two-box union β (33) is a smooth basis set.*

Proof: We will show that the conditions (a) and (b) of Theorem 5.1.1 hold for β .

First suppose that $c_j^{\mathbf{d}}$ is a non-standard arrow; we must show that $c_j^{\mathbf{d}}$ is translation-equivalent to 0. We may assume, by Lemma 4.1.2, that $c_j^{\mathbf{d}}$ is a minimal arrow; that is, its tail $\mathbf{x}^{\mathbf{d}}$ is a minimal generator of I_β . Since every arrow with tail a corner monomial $x_i^{w_i}$ is standard, Lemma 11.2.2 implies that

$$\mathbf{x}^{\mathbf{d}} = x_j^{w_{j,2}} x_k^{w_{k,1}}, \quad x_j \in V_1, \quad x_k \in V_2.$$

Without loss of generality, suppose that the head $\mathbf{x}^{\mathbf{j}} \in \mathcal{B}_1$. Then we can translate the arrow in the direction of increasing x_j -degree to reach $c_{j_1}^{\mathbf{d}_1}$, where

$$\begin{aligned} x_j\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) &= w_j - 1 = w_{j,1} - 1 \quad \text{and} \\ x_i\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) &= x_i\text{-degree}(\mathbf{x}^{\mathbf{j}}), \quad \text{for } i \neq j. \end{aligned}$$

Then

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{d}_1}) > x_j\text{-degree}(\mathbf{x}^{\mathbf{j}_1}) \Rightarrow x_j\text{-degree}(\mathbf{x}^{\mathbf{d}_1}) \geq w_j,$$

so we may translate $c_{j_1}^{\mathbf{d}_1}$ by degree-decreasing steps so that its tail approaches the corner monomial $x_j^{w_j}$. The head must eventually exit the first orthant, since a non-standard arrow cannot have a corner monomial as its tail. Therefore, $c_j^{\mathbf{d}} \sim 0$, and condition (a) holds.

Now let $\mathcal{S}$ be a standard bunch of arrows for β , and $c_j^{\mathbf{d}} \not\sim 0$ be a standard arrow for β . We must show that there exists an arrow $c_{j'}^{\mathbf{d}'} \in \mathcal{S}$ such that $c_j^{\mathbf{d}} \sim c_{j'}^{\mathbf{d}'}$. We may assume that $c_j^{\mathbf{d}}$ is a minimal x_i -standard arrow that cannot be advanced, by assertion (c) of Lemma 4.2.1. If the tail $\mathbf{x}^{\mathbf{d}}$ is the corner monomial $x_i^{w_i}$, we are done, since then

$$c_j^{\mathbf{d}} \in \mathcal{S}(i) \subseteq \mathcal{S}$$

by Corollary 4.6.2. Otherwise, by Lemma 11.2.2, we have that

$$\mathbf{x}^{\mathbf{d}} = x_j^{w_{j,2}} x_k^{w_{k,1}}, \quad x_j \in V_1, \quad x_k \in V_2,$$

and we may assume without loss of generality that $x_i = x_j$. Let v denote the x_j -offset of $c_j^{\mathbf{d}}$, and let $c_j^{\mathbf{d}'}$ be the unique arrow in $\mathcal{S}(j)$ such that

$$x_j\text{-degree}(\mathbf{x}^{\mathbf{j}'}) = x_j\text{-degree}(\mathbf{x}^{\mathbf{j}}) \quad \text{and} \quad x_j\text{-offset}(c_j^{\mathbf{d}'}) = v;$$

the existence of $c_j^{\mathbf{d}'}$ is guaranteed by Corollary 4.6.1. The tail $\mathbf{x}^{\mathbf{d}'}$ is a minimal generator of I_β that is divisible by x_j . We proceed to show that $c_j^{\mathbf{d}} \sim c_j^{\mathbf{d}'}$.

If $\mathbf{x}^{\mathbf{d}'} = x_j^{w_j}$, then one sees easily that

$$\mathbf{x}^{\mathbf{j}'} = \mathbf{x}^{\mathbf{j}} / x_k^{w_{k,1}};$$

it is then apparent that $c_j^{\mathbf{d}'}$ can be translated $w_{k,1}$ steps in the direction of increasing x_k -degree to reach an arrow $c_j^{\mathbf{d}_1'}$, whose tail $\mathbf{x}^{\mathbf{d}_1'}$ is divisible by $\mathbf{x}^{\mathbf{d}}$; therefore, $c_j^{\mathbf{d}_1'}$ and $c_j^{\mathbf{d}'}$ can be advanced, which is a contradiction. Hence, Lemma 11.2.1 yields that

$$\mathbf{x}^{\mathbf{d}'} = x_j^{w_{j,2}} x_{k'}^{w_{k',1}}, \quad x_{k'} \in V_2.$$

We now know that $c_j^{\mathbf{d}}$ and $c_j^{\mathbf{d}'}$ have the same length, x_j -offset, and x_j -height = $w_{j,2}$. By Lemma 11.3.1, we know that

$$\mathbf{x}^{\mathbf{j}}, \mathbf{x}^{\mathbf{j}'} \in \mathcal{B}_2 \Rightarrow \text{lcm}(\mathbf{x}^{\mathbf{j}}, \mathbf{x}^{\mathbf{j}'}) = \mathbf{x}^{\mathbf{j}^*} \in \mathcal{B}_2 \subseteq \beta,$$

where the implication uses Lemma 6.5.2. It is now clear that $c_j^{\mathbf{d}}$ and $c_j^{\mathbf{d}'}$ can each be translated by degree-increasing steps to the same arrow $c_j^{\mathbf{d}^*}$; whence, $c_j^{\mathbf{d}} \sim c_j^{\mathbf{d}'}$, as desired. Therefore, condition (b) holds, and the proof is complete. $\square$

Remark 11.3.3. Note that *three*-box unions need not be smooth; for example, the left-hand basis set illustrated in Figure 9 is a three-box union, but is not smooth, since a non-standard rigid arrow exists.

11.4. Example: $\beta = \{1, x_1, x_2, x_1x_2, x_3, x_4, x_3x_4\}$. This basis set β is the two-box union

$$\beta = \mathcal{B}(2, 2, 1, 1) \cup \mathcal{B}(1, 1, 2, 2) \subseteq \mathbf{k}[x_1, x_2, x_3, x_4].$$

Therefore, by Theorem 11.3.2, β is a smooth basis set. However, it is clear that β is not a compound box, so the characterization of smooth basis sets in three variables given by Theorem 10.3.1 does not extend to higher dimensions.

REFERENCES

1. J. Fogarty, *Algebraic families on an algebraic surface*, Am. J. Math. **90** (1968), 511–521.
2. A. Geramita, D. Gregory, and L. Roberts, *Monomial ideals and points in projective space*, J. Pure Appl. Algebra **40** (1986), 33–62.
3. M. Gordan, *Les invariants des formes binaires*, J. Math Pures et Appl. (Liouville's J.) **6** (1900), 141–156.
4. A. Grothendieck, *Techniques de construction et théorèmes d'existence en géométrie algébrique, IV: Les schemas de Hilbert*, Sémin. Bourbaki 221, IHP, Paris, 1961.
5. T. S. Gustavsen, D. Laksov, and R. M. Skjelnes, *An elementary, explicit, proof of the existence of Hilbert schemes of points*, arXiv:math.AG/0506161, v1, 9 June 2005.

6. M. Haiman, *t, q-Catalan numbers and the Hilbert scheme*, Discrete Math. **193** (1998), no. 1-3, 201–224.
7. M. Haiman and B. Sturmfels, *Multigraded Hilbert schemes*, J. Algebraic Geom. **13** (2004), no. 4, 725–769.
8. R. Hartshorne, *Connectedness of the Hilbert scheme*, Publ. Math. Inst. Hautes Études Sci. **29** (1966), 263–304.
9. M. Huibregtse, *An elementary construction of the multigraded Hilbert scheme of points*, to appear in Pac. J. Math.
10. A. Iarrobino, *Compressed algebras and components of the punctual Hilbert scheme*, Algebraic Geometry Sitges (Barcelona) 1983 (E. Casas-Alvero, G. E. Welters, and S. Xambó-Descamps, eds.), Lecture Notes in Math., vol. 1124, Springer-Verlag, 1985, pp. 146–165.
11. R. Notari and M. L. Spreafico, *A stratification of Hilbert schemes by initial ideals and applications*, Manuscripta Math. **101** (2000), no. 4, 429–448.
12. A. Reeves and M. Stillman, *Smoothness of the lexicographic point*, J. Algebraic Geom. **6** (1997), 235–246.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, SKIDMORE COLLEGE, SARATOGA SPRINGS, NEW YORK 12866

E-mail address: mhuibreg@skidmore.edu